

SOLVMANIFOLDS AND NONCOMMUTATIVE TORI WITH REAL MULTIPLICATION

MATILDE MARCOLLI

ABSTRACT. We prove that the Shimizu L -function of a real quadratic field is obtained from a (Lorentzian) spectral triple on a noncommutative torus with real multiplication, as an adiabatic limit of the Dirac operator on a 3-dimensional solvmanifold. The Dirac operator on this 3-dimensional geometry gives, via the Connes–Landi isospectral deformations, a spectral triple for the noncommutative tori obtained by deforming the fiber tori to noncommutative spaces. The 3-dimensional solvmanifold is the homotopy quotient in the sense of Baum–Connes of the noncommutative space obtained as the crossed product of the noncommutative torus by the action of the units of the real quadratic field. This noncommutative space is identified with the twisted group C^* -algebra of the fundamental group of the 3-manifold. The twisting can be interpreted as the cocycle arising from a magnetic field, as in the theory of the quantum Hall effect. We prove a twisted index theorem that computes the range of the trace on the K -theory of this noncommutative space and gives an estimate on the gaps in the spectrum of the associated Harper operator.

CONTENTS

| | |
|-----------------------------------------------------------------------|----|
| 1. Introduction | 2 |
| 2. 3-dimensional solvmanifolds and real quadratic fields | 4 |
| 2.1. Semidirect products and solvmanifolds | 4 |
| 2.2. The topology of the 3-manifold X_ϵ | 5 |
| 2.3. Action on \mathbb{Z}^2 | 6 |
| 2.4. Solvmanifold and Hecke’s lift of geodesics | 6 |
| 3. Actions on noncommutative tori with real multiplication | 7 |
| 4. Twisted group algebras and the magnetic Laplacian | 8 |
| 4.1. Twisted group algebras | 9 |
| 4.2. The noncommutative tori as twisted group algebras | 9 |
| 4.3. Twisted group algebra of $S(\Lambda, V)$ | 10 |
| 4.4. The magnetic Laplacian | 11 |
| 4.5. Discretized electron–ion interaction and Harper operators | 12 |
| 4.6. Harper operators for noncommutative tori and for $S(\Lambda, V)$ | 13 |
| 4.7. Spectral theory and K -theory | 13 |
| 5. Homotopy quotient and the Baum–Connes conjecture | 14 |
| 5.1. K -theory of $C^*(S(\Lambda, V), \tilde{\sigma})$ | 14 |
| 5.2. K -theory and the twist | 15 |
| 5.3. Thom isomorphism, homotopy quotients and Baum–Connes | 16 |

| | | |
|------|----------------------------------------------------------------|----|
| 6. | Twisted index theorem, K -theory, and the range of the trace | 16 |
| 6.1. | Spectral flow and odd Chern character | 17 |
| 6.2. | Twisted index theorem | 18 |
| 6.3. | Range of the trace | 19 |
| 7. | Isospectral deformations and spectral triples | 21 |
| 7.1. | The Connes–Landi isospectral deformations | 21 |
| 7.2. | Noncommutative solvmanifolds | 23 |
| 7.3. | Unitary equivalences | 25 |
| 7.4. | Differential operators on noncommutative tori | 27 |
| 8. | Shimizu L -function and Lorentzian geometry | 28 |
| 8.1. | Arithmetic Krein spaces | 30 |
| 8.2. | Lorentzian spectral triples over real quadratic fields | 31 |
| 8.3. | Arithmetic twisted group algebras | 32 |
| 8.4. | Lorentzian Dirac operator | 34 |
| 8.5. | Eta function and 3-dimensional geometry | 36 |
| 8.6. | The residue | 38 |
| | References | 39 |

1. INTRODUCTION

In the 1970s Hirzebruch formulated a conjecture, *cf.* [15], on the topological interpretation of certain special values of L -functions of totally real fields in terms of signature defects. The conjecture was proved in the early '80s by Atiyah–Donnelly–Singer [2] and by Müller [23]. Hirzebruch's conjecture played an important role in the development of the Atiyah–Patodi–Singer index theorem [3], which in turn is a key ingredient in the proof [2] of the conjecture, extending the Hirzebruch–Riemann–Roch theorem to manifolds with boundary and relating the signature defect to the eta invariant. Geometrically, the link of an isolated singularity of the Hilbert modular variety associated to a totally real number field is given by a $(4k - 1)$ -dimensional solvmanifold. The signature of the Hilbert modular variety is then computed by the APS theorem applied to the resulting manifold with boundary and the signature defects are computed by the eta invariant of the solvmanifold. The main step in the proof of [2] then consists of separating out the eta function of the signature operator on the solvmanifold into a part that recovers the Shimizu L -function of the totally real field and a residual part, which is shown not to contribute to the eta invariant.

We concentrate here on the simplest case, that of real quadratic fields, and we consider the question of whether the Shimizu L -function can be related in a similar way to a spectral geometry (in the sense of Connes' notion of spectral triples [11]) on a noncommutative torus with real multiplication by the same real quadratic field.

The noncommutative tori associated to quadratic irrationalities have been extensively studied by Manin in [17] and subsequently by several authors. They have the special property of “real multiplication”, derived from the presence of non-trivial self Morita equivalences. It is argued in [17] that the noncommutative tori with real

multiplication should play a role for real quadratic field parallel to the the theory of elliptic curves with complex multiplication in the case of imaginary quadratic fields. This makes it an interesting problem to study the geometric properties of this particular class of noncommutative spaces, and their relation to the arithmetic of real quadratic fields.

We show in §5 that there is a close relation between the 3-dimensional solvmanifold and the noncommutative torus with real multiplication. Namely, we prove that the first is the homotopy quotient, in the sense of Baum–Connes, of the noncommutative space given by the quotient of the latter by the action of the infinite group of units, *cf.* §3. We also show that the 3-manifold can be identified with the pullback of the universal family of elliptic curves along a closed geodesic in the modular curve.

This interpretation as the homotopy quotient of a noncommutative space provides a geometric setting analogous to the one developed in the noncommutative geometry models of the quantum Hall effect [5], where the presence of a magnetic field makes the Brillouin zone of the lattice into a noncommutative torus. Here, the 3-dimensional solvmanifold is similarly related to a noncommutative space whose algebra of coordinates is the crossed product of the algebra of the noncommutative torus by the action of the units. This is obtained by twisting the group ring of the fundamental group of the solvmanifold by a cocycle, defined in terms of a magnetic potential. The noncommutative space is the resulting twisted group C^* -algebra, *cf.* §4. As in the case of the quantum Hall effect, and in the noncommutative Bloch theory of electron-ion interactions, one obtains in §6 information on the spectral theory of the corresponding magnetic Laplacian by computing the range of the trace on the K -theory of the twisted group C^* -algebra. We prove a twisted index theorem which we use to compute the range of the trace using a spectral flow computation and the Baum–Connes conjecture, which is known to hold for the fundamental group of the 3-dimensional solvmanifold.

This way of passing from the 3-dimensional solvmanifold to the associated noncommutative space is obtained in two steps. Viewing the 3-manifold as a fibration of 2-dimensional tori over the circle, one first replaces the fiber tori by noncommutative tori and then the mapping torus by the dual action of the units. We show in §7 that the first step can be seen as a case of the Connes–Landi isospectral deformations [12]. In particular, we prove that the Dirac operator on the 3-manifold induces in this way a Dirac operator on the noncommutative torus with real multiplication. A unitary equivalence as the one considered in [2] then factors this Dirac operator into a product of two operators, one of which has spectrum given by the norms $N(\lambda)$ of the lattice points λ and recovers the Shimizu L -function. In §7.4 we show how an adiabatic limit relates the Dirac operator on the 3-manifold to known differential operators on the noncommutative torus, obtained by considering the derivations along the leaves of the Kronecker foliations associated to the Galois conjugate elements θ and θ' in the real quadratic field $\mathbb{K} = \mathbb{Q}(\theta)$.

Finally we prove in §8 that the norms $N(\lambda)$ define the momenta of a *Lorentzian* Dirac operator on the noncommutative torus with real multiplication. The theory of spectral triples in Lorentzian signature is at present still under active development

and this provides a natural example where the arithmetic structure dictates how the Lorentzian geometry should be treated in the noncommutative context. We develop a framework for Lorentzian spectral triples over real quadratic fields, where the Galois involution of \mathbb{K} provides a canonical choice of a Krein involution. In particular, we prove that, in passing from the indefinite Lorentzian geometry defined by the quadratic form given by the norm to the associated real Hilbert space, one can resolve the infinite multiplicities in the spectrum of the Dirac operator arising from the presence of a non-compact group of symmetries (the units of the real quadratic field). We show that the resulting operator on the real Hilbert space has the same eta function as the one coming from the adiabatic limit of the 3-dimensional geometry as in §7.4. This eta function is the product of the Shimizu L -function of the real quadratic field by a term that only depends on the fundamental unit.

Acknowledgment. I am very grateful to Sir Michael Atiyah for asking the question this paper is attempting to answer. I thank Alain Connes, Yuri Manin, and Don Zagier for useful conversations. I also thank the Mittag Leffler Institute for hospitality and support, while part of this work was done. This research is partially supported by NSF grant DMS-0651925.

2. 3-DIMENSIONAL SOLVMANIFOLDS AND REAL QUADRATIC FIELDS

Let $\mathbb{K} = \mathbb{Q}(\sqrt{d})$ be a real quadratic field and let $\iota_i : \mathbb{K} \hookrightarrow \mathbb{R}$, for $i = 1, 2$, be its two real embeddings. We let $L \subset \mathbb{K}$ be a lattice, with U_L^+ the group of totally positive units preserving L ,

$$(2.1) \quad U_L^+ = \{u \in O_{\mathbb{K}}^* \mid uL \subset L, \iota_i(u) \in \mathbb{R}_+^*\}.$$

We denote by ϵ_L a generator, so that $U_L^+ = \epsilon_L^{\mathbb{Z}}$. In the case where $L = O_{\mathbb{K}}$, the ring of integers of \mathbb{K} , then the generator $\epsilon = \epsilon_L$ is a fundamental unit. We consider the embedding of L in \mathbb{R}^2 given by the mapping

$$(2.2) \quad L \ni \ell \mapsto (\iota_1(\ell), \iota_2(\ell)) \in \mathbb{R}^2.$$

We denote the range by $\Lambda = (\iota_1, \iota_2)(L)$. This is a lattice in \mathbb{R}^2 . The action of U_L^+ extends to an action on Λ by

$$(2.3) \quad \lambda = (\iota_1(\ell), \iota_2(\ell)) \mapsto (\epsilon \iota_1(\ell), \epsilon' \iota_2(\ell)) = (\epsilon \iota_1(\ell), \epsilon^{-1} \iota_2(\ell)).$$

2.1. Semidirect products and solvmanifolds. Let us denote by V either the group U_L^+ or a finite index subgroup thereof. As in [2], we consider the crossed product

$$(2.4) \quad S(\Lambda, V) = \Lambda \rtimes_{\epsilon} V,$$

where the action of $V = \epsilon^{\mathbb{Z}}$ on Λ is induced by the action by multiplication on L . As shown in [2], these are discrete subgroups of the solvable Lie group

$$(2.5) \quad S(\mathbb{R}^2, \mathbb{R}) = \mathbb{R}^2 \rtimes \mathbb{R},$$

with the action of \mathbb{R} on \mathbb{R}^2 by the one parameter subgroup $\Theta_t(x, y) = (e^t x, e^{-t} y)$ of $\mathrm{SL}_2(\mathbb{R})$. For $\epsilon > 1$ and $\epsilon' = \epsilon^{-1} < 1$, the action of V on Λ is then generated by

$$(2.6) \quad A_\epsilon = \begin{pmatrix} \epsilon & 0 \\ 0 & \epsilon' \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R}).$$

We also consider the 3-dimensional solvmanifold obtained as the quotient

$$(2.7) \quad X_\epsilon = S(\Lambda, V) \backslash S(\mathbb{R}^2, \mathbb{R}),$$

with $\pi_1(X_\epsilon) = S(\Lambda, V)$.

2.2. The topology of the 3-manifold X_ϵ . It is well known [2] that the 3-manifold X_ϵ of (2.7) is a fibration over the circle S^1 with fibers that are 2-tori and with monodromy given by the matrix A_ϵ of (2.6).

Lemma 2.1. *The manifold X_ϵ has first homology*

$$(2.8) \quad H_1(X_\epsilon, \mathbb{Z}) = \Lambda / (1 - A_\epsilon)\Lambda \oplus \mathbb{Z}.$$

Proof. The fundamental group is $\pi_1(X_\epsilon) = S(\Lambda, V)$. Consider the surjective map

$$(2.9) \quad \pi : S(\Lambda, V) \rightarrow \Lambda / (1 - A_\epsilon)\Lambda \oplus \mathbb{Z}, \quad \pi(\lambda, n) = (\lambda \bmod (1 - A_\epsilon)\Lambda, n).$$

By writing

$$A_\epsilon^n(\lambda') = \lambda' - (1 - A_\epsilon)(\lambda' + A_\epsilon(\lambda') + \cdots + A_\epsilon^{n-1}(\lambda'))$$

one sees that $\lambda + A_\epsilon^n(\lambda') = \lambda + \lambda'$ modulo $(1 - A_\epsilon)\Lambda$, so that $\pi(\lambda + A_\epsilon^n \lambda', n + n') = \pi(\lambda, n) + \pi(\lambda', n')$. Since commutators in $S(\Lambda, V)$ are of the form

$$(\lambda, n)(\lambda', n')(\lambda, n)^{-1}(\lambda', n')^{-1} = ((1 - A_\epsilon^{n'})\lambda - (1 - A_\epsilon^n)\lambda', 0),$$

we see that the homomorphism (2.9) has $\mathrm{Ker}(\pi) = [\pi_1(X_\epsilon), \pi_1(X_\epsilon)]$. \square

Corollary 2.2. *The compact 3-manifold X_ϵ has cohomology*

$$(2.10) \quad H^{\mathrm{even}}(X_\epsilon, \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathrm{Coker}(1 - A_\epsilon), \quad H^{\mathrm{odd}}(X_\epsilon, \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}.$$

Proof. By Poincaré duality we have

$$(2.11) \quad H^2(X_\epsilon, \mathbb{Z}) \cong H_1(X_\epsilon, \mathbb{Z}) \cong \mathbb{Z} \oplus \Lambda / (1 - A_\epsilon)\Lambda,$$

and $H^1(X_\epsilon, \mathbb{Z}) = \mathrm{Hom}(H_1(X_\epsilon, \mathbb{Z}), \mathbb{Z}) = \mathbb{Z} \oplus \mathrm{Hom}(\Lambda / (1 - A_\epsilon)\Lambda, \mathbb{Z})$, so that

$$(2.12) \quad H_2(X_\epsilon, \mathbb{Z}) \cong H^1(X_\epsilon, \mathbb{Z}) \cong \mathbb{Z} \oplus \mathrm{Hom}(\Lambda / (1 - A_\epsilon)\Lambda, \mathbb{Z}).$$

We have $\mathrm{Ker}(1 - A_\epsilon) = 0$, while $\mathrm{Coker}(1 - A_\epsilon)$ is torsion, so that we obtain

$$(2.13) \quad \begin{aligned} H^0(X_\epsilon, \mathbb{Z}) &= \mathbb{Z} & H^1(X_\epsilon, \mathbb{Z}) &= \mathbb{Z} \\ H^3(X_\epsilon, \mathbb{Z}) &= \mathbb{Z} & H^2(X_\epsilon, \mathbb{Z}) &= \mathbb{Z} \oplus \mathrm{Coker}(1 - A_\epsilon) \end{aligned}$$

\square

2.3. Action on \mathbb{Z}^2 . We recall the following description of the action of A_ϵ on Λ , which will be useful in the following, where we use twisted group C^* -algebras to describe noncommutative tori.

Lemma 2.3. *In the basis $\{1, \theta\}$ of $\iota_1(L) \subset \mathbb{R}$, the action of the group $V = \epsilon^\mathbb{Z}$ is generated by the matrix*

$$(2.14) \quad \varphi_\epsilon = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$

with $\epsilon = a + b\theta$ and $\epsilon\theta = c + d\theta$. The conjugate elements $1/\theta$ and $1/\theta'$ are the fixed points of $\varphi_\epsilon \in \mathrm{SL}_2(\mathbb{Z})$ acting on $\mathbb{P}^1(\mathbb{R})$ by fractional linear transformations.

Proof. As we have seen in (2.3), the action of V on Λ is given by

$$A_\epsilon : (n + m\theta, n + m\theta') \mapsto (\epsilon(n + m\theta), \epsilon'(n + m\theta'))$$

with $\epsilon\epsilon' = 1$. In particular, for $m = 0$ and $n = 1$ this gives $\epsilon \in \iota_1(L)$ and $\epsilon' \in \iota_2(L)$. Thus, we can write $\epsilon = a + b\theta$, for two integers $a, b \in \mathbb{Z}$. Similarly, the element $\epsilon\theta$ can be written in the form $\epsilon\theta = c + d\theta$. Thus, the action of A_ϵ^k on Λ can be described equivalently as

$$(2.15) \quad (n, m) \mapsto (n, m)\varphi_\epsilon^k, \quad \text{with } \varphi_\epsilon = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$$

The second statement follows immediately since

$$\theta^{-1} = \epsilon/(\epsilon\theta) = (a\theta^{-1} + b)/(c\theta^{-1} + d).$$

□

We obtain in this way two corresponding identifications $S(\Lambda, V) = \mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}$, by mapping (λ, ϵ^k) to either $(\lambda_1 = n + m\theta, k)$ or $(\lambda_2 = n + m\theta', -k)$.

2.4. Solvmanifold and Hecke's lift of geodesics. For $\Gamma = \mathrm{SL}_2(\mathbb{Z})$ and $X_\Gamma = \Gamma \backslash \mathbb{H}$ the modular curve, let $\mathcal{U}_\Gamma \rightarrow X_\Gamma$ denote the universal family of elliptic curves over the modular curve, where the fiber over $\tau \in X_\Gamma$ of \mathcal{U}_Γ is the isomorphism class of the elliptic curve $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$.

Suppose given a lattice L in a real quadratic field \mathbb{K} and let $\{1, \theta\}$ be a basis for L , with θ' the Galois conjugate of θ under the Galois involution of \mathbb{K} over \mathbb{Q} .

We denote by $\gamma_{\theta, \theta'}$ the infinite geodesic in the hyperbolic plane \mathbb{H} with endpoints $\theta, \theta' \in \mathbb{P}^1(\mathbb{R})$. This defines a closed geodesic in the quotient X_Γ of length $\log \epsilon$, for $\epsilon > 1$ the generator of $V = U_L^+ = \epsilon^\mathbb{Z}$. We denote the closed geodesic by $\bar{\gamma}_{\theta, \theta'}$.

Consider the restriction of the universal family \mathcal{U}_Γ to the closed geodesic $\bar{\gamma}_{\theta, \theta'}$. Via the parameterization of the closed geodesic by a circle S^1 of length $\log \epsilon$, we can consider the pullback to the parameterizing S^1 of \mathcal{U}_Γ . We obtain in this way a real 3-dimensional manifold, which we denote $\mathcal{U}_{\theta, \theta'}$. This is topologically a 3-manifold that fibers over a circle, with fibers T^2 . We consider it endowed with the metric that is the product of the geodesic length and the flat metric on T^2 . We then obtain the following result.

Lemma 2.4. *The solvmanifold $S(\Lambda, V)$ is isometrically equivalent to $\mathcal{U}_{\theta, \theta'}$.*

Proof. We recall the following procedure of Hecke to lift closed geodesics to the space of lattices (cf. Manin [17], §1.8.2). Given a lattice L in a real quadratic field \mathbb{K} , with $\ell \rightarrow \ell'$ the Galois involution, one sets

$$(2.16) \quad \Lambda_t(L) := \{z \in \mathbb{H} \mid z = z(\ell, t) = \ell e^t + i\ell' e^{-t} \mid \ell \in L\}.$$

This defines, for all $t \in \mathbb{R}$ a lattice $\Lambda_t \subset \mathbb{C}$. The action of $V = \epsilon^{\mathbb{Z}}$ is of the form (cf. Lemma 1.8.3 of [17])

$$(2.17) \quad z(\ell, t) \mapsto \epsilon \ell e^t + i\epsilon' \ell' e^{-t} = z(\ell, t + \log \epsilon).$$

In particular (see again [17] Lemma 1.8.3), for $\{1, \theta\}$ a basis of L , the lattice $\Lambda_t(L)$ is generated by $\{1, \tau_t\}$ where τ_t runs over the geodesic $\ell_{\theta, \theta'} \subset \mathbb{H}$, for $t \in \mathbb{R}$. Thus, we can identify the 3-manifold $\mathcal{U}_{\theta, \theta'}$ with the fibration over a circle of length $\log \epsilon$, with fiber $E_{\tau_t} = \mathbb{C}/\Lambda_t(L)$.

On the other hand, the 3-manifold $S(\Lambda, V)$ is a fibrations of tori over the circle

$$(2.18) \quad T^2 \rightarrow S(\Lambda, V) \rightarrow S^1,$$

where the base S^1 is a circle of length $\log \epsilon$ and the fiber over $t \in S^1$ is given by the 2-torus

$$(2.19) \quad T_t^2 = \mathbb{R}^2/\Lambda_t,$$

with $\Lambda_t = \Theta_t(\Lambda)$, for $\Theta_t(x, y) = (e^t x, e^{-t} y)$. This proves the result. \square

3. ACTIONS ON NONCOMMUTATIVE TORI WITH REAL MULTIPLICATION

The noncommutative torus \mathcal{A}_θ of modulus $\theta \in \mathbb{R} \setminus \mathbb{Q}$ is the noncommutative space described, at the topological level, by the irrational rotation C^* -algebra, that is, the universal C^* -algebra generated by two unitaries U, V with the commutation relation $VU = e^{2\pi i \theta} UV$. It has a smooth structure given by the smooth subalgebra of series $\sum_{n,m} a_{n,m} U^n V^m$ with rapidly decaying coefficients (cf. [9]).

It is a well known result ([9], [26]) that the algebras \mathcal{A}_{θ_1} and \mathcal{A}_{θ_2} are Morita equivalent whenever there exists an element $g \in \text{SL}_2(\mathbb{Z})$ acting on \mathbb{R} by fractional linear transformations, such that $\theta_1 = g\theta_2$. In the following we concentrate on the case where the irrational number θ is a quadratic irrationality in a real quadratic field $\mathbb{K} = \mathbb{Q}(\theta)$. These are the noncommutative tori with real multiplication considered in [17]. We let L be the lattice in \mathbb{K} with $\iota_1(L) = \mathbb{Z} + \mathbb{Z}\theta$ and $\iota_2(L) = \mathbb{Z} + \mathbb{Z}\theta'$. As before, we denote by Λ the corresponding lattice in \mathbb{R}^2 .

The C^* -algebra of the noncommutative torus \mathcal{A}_θ described above can be equivalently described as the crossed product

$$(3.1) \quad \mathcal{A}_\theta = C(S^1) \rtimes_\theta \mathbb{Z},$$

where the action of \mathbb{Z} on S^1 is by the irrational rotation by $\exp(2\pi i \theta)$. Up to Morita equivalence, one can replace $C(S^1)$ by the crossed product $C_0(\mathbb{R}) \rtimes \mathbb{Z}$, and one obtains a Morita equivalent description of the noncommutative torus as

$$(3.2) \quad C_0(\mathbb{R}) \rtimes_\theta \mathbb{Z}^2 = C_0(\mathbb{R}) \rtimes (\mathbb{Z} + \mathbb{Z}\theta).$$

In the case we are considering, of real quadratic fields, we can regard the noncommutative torus with real multiplication associated to a lattice $L \subset \mathbb{K}$ as described by the algebras

$$(3.3) \quad \mathbb{T}_{\Lambda,1} := C_0(\mathbb{R}) \rtimes \iota_1(L) \quad \mathbb{T}_{\Lambda,2} := C_0(\mathbb{R}) \rtimes \iota_2(L).$$

These algebras can be described as follows. They are C^* -algebras generated by elements of the form fU_λ , with $f \in C_0(\mathbb{R})$ and $\lambda \in \Lambda$, with the product

$$fU_\lambda hU_\eta = fU_{\lambda,i}(h)U_{\lambda+\eta}, \quad \text{where} \quad U_{\lambda,i}(h)(x) = h(x + \iota_i(\lambda)), \quad i = 1, 2.$$

The group $V = \epsilon^\mathbb{Z}$ of units acts as symmetries of the noncommutative tori $\mathbb{T}_{\Lambda,i}$ as follows.

Lemma 3.1. *For $k \in \mathbb{Z}$ and $fU_\lambda \in \mathbb{T}_{\Lambda,i}$, set $v_\epsilon^k(f)(x) := f(\epsilon^k x)$ and*

$$(3.4) \quad v_1^k(fU_\lambda) = v_\epsilon^k(f)U_{A_\epsilon^k(\lambda)}, \quad \text{and} \quad v_2^k(fU_\lambda) = v_{\epsilon'}^k(f)U_{A_{\epsilon'}^k(\lambda)}.$$

This defines actions $v_i : V \rightarrow \text{Aut}(\mathbb{T}_{\Lambda,i})$.

Proof. The result follows directly from

$$v_i^k(U_\lambda(h))(x) = U_{A_\epsilon^k(\lambda)}(v_i^k(h))(x) = \begin{cases} h(\epsilon^k(x + n + m\theta)) & i = 1 \\ h(\epsilon^{-k}(x + n + m\theta')) & i = 2 \end{cases}$$

which implies that

$$v_\epsilon^k(fU_\lambda hU_\eta) = v_i^k(f)v_i^k(U_\lambda(h))U_{A_\epsilon^k(\lambda+\eta)} = v_i^k(fU_\lambda)v_i^k(hU_\eta).$$

□

It is customary, in noncommutative geometry, to replace quotients by crossed product algebras. In this case, the quotient of the noncommutative tori $\mathbb{T}_{\Lambda,i}$ by the action of V is described by the crossed product algebra

$$(3.5) \quad \mathbb{T}_{\Lambda,V,i} := \mathbb{T}_{\Lambda,i} \rtimes_{v_i} V,$$

which we can view equivalently as the crossed product

$$(3.6) \quad \mathbb{T}_{\Lambda,V,i} := C_0(\mathbb{R}) \rtimes_i S(\Lambda, V),$$

for the actions of $S(\Lambda, V)$ on $C_0(\mathbb{R})$ of the form

$$(3.7) \quad U_{(\lambda,k)}f(x) = f(\epsilon^k(x + n + m\theta)) \quad \text{or} \quad U_{(\lambda,k)}f(x) = f(\epsilon^{-k}(x + n + m\theta')).$$

4. TWISTED GROUP ALGEBRAS AND THE MAGNETIC LAPLACIAN

Another equivalent description of the algebra \mathcal{A}_θ of the noncommutative torus is as twisted group C^* -algebra. This played an important role in the context of the noncommutative geometry model of the integer quantum Hall effect (see [5]).

We recall briefly the definition and properties of twisted group C^* -algebras, as this will be useful in the following. For a similar overview and applications to the case of Fuchsian groups see [20].

4.1. Twisted group algebras. Let Γ be a finitely generated discrete group, and let $\sigma : \Gamma \times \Gamma \rightarrow U(1)$ be a multiplier, that is, a 2-cocycle satisfying the cocycle property

$$(4.1) \quad \sigma(\gamma_1, \gamma_2)\sigma(\gamma_1\gamma_2, \gamma_3) = \sigma(\gamma_1, \gamma_2\gamma_3)\sigma(\gamma_2, \gamma_3),$$

with $\sigma(\gamma, 1) = \sigma(1, \gamma) = 1$.

Consider then the Hilbert space $\ell^2(\Gamma)$ and the left/right σ -regular representations of Γ given by

$$(4.2) \quad L_\gamma^\sigma f(\gamma') = f(\gamma^{-1}\gamma')\sigma(\gamma, \gamma^{-1}\gamma'), \quad R_\gamma^\sigma f(\gamma') = f(\gamma'\gamma)\sigma(\gamma', \gamma).$$

They satisfy the relations

$$(4.3) \quad L_\gamma^\sigma L_{\gamma'}^\sigma = \sigma(\gamma, \gamma')L_{\gamma\gamma'}^\sigma, \quad R_\gamma^\sigma R_{\gamma'}^\sigma = \sigma(\gamma, \gamma')R_{\gamma\gamma'}^\sigma.$$

Moreover the left σ -regular representation commutes with the right $\bar{\sigma}$ -regular representation, with $\bar{\sigma}$ the conjugate multiplier. The algebra generated by the R_γ^σ is the twisted group ring $\mathbb{C}(\Gamma, \sigma)$. Its norm closure is the (reduced) twisted group C^* -algebra $C_r^*(\Gamma, \sigma)$.

4.2. The noncommutative tori as twisted group algebras. One identifies the C^* -algebra \mathcal{A}_θ of the noncommutative torus with the reduced twisted group C^* -algebra $C_r^*(\mathbb{Z}^2, \sigma)$ in the following way. Let σ be a cocycle of the form

$$(4.4) \quad \sigma((n, m), (n', m')) := \exp(-2\pi i(\xi_1 nm' + \xi_2 mn')).$$

Then the operators $U = R_{(0,1)}^\sigma$ and $V = R_{(1,0)}^\sigma$ acting by

$$Uf(n, m) = e^{-2\pi i\xi_2 n} f(n, m+1), \quad Vf(n, m) = e^{-2\pi i\xi_1 m} f(n+1, m)$$

that generate the algebra $C_r^*(\mathbb{Z}^2, \sigma)$ satisfy the commutation relation

$$UV = e^{2\pi i\theta} VU, \quad \text{with } \theta = \xi_2 - \xi_1.$$

Notice that different choices of ξ_1, ξ_2 with the same $\theta = \xi_2 - \xi_1$ yield the same algebra \mathcal{A}_θ . This gives us the freedom to choose the ξ_i according to the following result.

Lemma 4.1. *A cocycle σ of the form (4.4) has the property that*

$$(4.5) \quad \sigma((n, m), (n', m')) = \sigma((n, m)\varphi, (n', m')\varphi), \quad \forall \varphi = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z})$$

if and only if $\xi_2 = -\xi_1$.

Proof. We see that $\sigma((n, m)\varphi, (n', m')\varphi)$ is of the form

$$\exp(-2\pi i((\xi_1 + \xi_2)(abnn' + cdmm') + (\xi_1 cb + \xi_2 ad)mn' + (\xi_1 ad + \xi_2 cb)nm')).$$

□

Thus, in the following we will assume that $\xi_2 = \theta/2 = -\xi_1$ in the choice of the cocycle σ of (4.4). We can then write σ in the form

$$\sigma_\theta((n, m), (k, r)) = \exp(\pi i\theta(nr - mk)) = \exp(\pi i\theta(n, m) \wedge (k, r)),$$

where we use the notation

$$(4.6) \quad (a, b) \wedge (c, d) = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

We then obtain the following identifications.

Corollary 4.2. *The noncommutative tori $\mathbb{T}_{\Lambda, i}$ are described by twisted group C^* -algebras*

$$(4.7) \quad \begin{aligned} \mathbb{T}_{\Lambda, 1} &= C^*(\mathbb{Z}^2, \sigma_\theta) = C^*(\Lambda, \sigma_{\theta(\theta' - \theta)^{-1}}), \\ \mathbb{T}_{\Lambda, 2} &= C^*(\mathbb{Z}^2, \sigma_{\theta'}) = C^*(\Lambda, \sigma_{\theta'(\theta' - \theta)^{-1}}). \end{aligned}$$

Proof. The expression

$$\sigma_u(\lambda, \eta) = \exp(\pi i u \lambda \wedge \eta)$$

defines a cocycle on Λ . For $\lambda = (n + m\theta, n + m\theta')$ and $\eta = (k + r\theta, k + r\theta')$, a direct calculation shows that

$$\sigma_\theta((n, m), (k, r)) = \sigma_u(\lambda, \eta), \quad \text{for } u = \theta(\theta' - \theta)^{-1}.$$

Thus, the generators $R_{(n, m)}^\sigma$ of $C^*(\mathbb{Z}^2, \sigma_\theta)$ with

$$R_{(n, m)}^\sigma R_{(k, r)}^\sigma = \sigma_\theta((n, m), (k, r)) R_{(n, m) + (k, r)}^\sigma$$

are identified with the generators R_λ^σ of $C^*(\Lambda, \sigma_{\theta(\theta' - \theta)^{-1}})$ with

$$R_\lambda^\sigma R_\eta^\sigma = \sigma_{\theta(\theta' - \theta)^{-1}}(\lambda, \eta) R_{\lambda + \eta}^\sigma.$$

The case of $\mathbb{T}_{\Lambda, 2}$ is analogous. □

4.3. Twisted group algebra of $S(\Lambda, V)$. We now show that the algebra $\mathcal{A}_\theta \rtimes V$, which we introduced in the previous section to describe the quotient of the noncommutative torus with real multiplication by the action of V , also admits a description in terms of twisted group C^* -algebras, for the group $S(\Lambda, V)$. First notice that the group $S(\Lambda, V)$ is amenable, so that the maximal and reduced group C^* -algebras coincide, $C_{max}^*(S(\Lambda, V)) \cong C_r^*(S(\Lambda, V))$, so that we can simply write $C^*(S(\Lambda, V))$, and $C^*(S(\Lambda, V), \tilde{\sigma})$ for the twisted case.

Lemma 4.3. *Let σ be a multiplier on \mathbb{Z}^2 of the form (4.4), with $\xi_2 = \theta/2 = -\xi_1$. Then the map $\tilde{\sigma} : S(\Lambda, V) \times S(\Lambda, V) \rightarrow U(1)$ of the form*

$$(4.8) \quad \tilde{\sigma}((n, m, k), (n', m', k')) := \sigma((n, m), (n', m')\varphi_\epsilon^k)$$

is a multiplier for $S(\Lambda, V)$, identified with the group $\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}$.

Proof. The cocycle condition for σ and the $\text{SL}_2(\mathbb{Z})$ -invariance $\sigma((n, m)\varphi, (n', m')\varphi) = \sigma((n, m), (n', m'))$ imply that $\tilde{\sigma}$ also satisfies the cocycle condition (4.1), since we have

$$\begin{aligned} &\sigma((n_1, m_1), (n_2, m_2)\varphi_\epsilon^{k_1})\sigma((n_1, m_1) + (n_2, m_2)\varphi_\epsilon^{k_1}, (n_3, m_3)\varphi_\epsilon^{k_1+k_2}) = \\ &\sigma((n_1, m_1), (n_2, m_2)\varphi_\epsilon^{k_1} + (n_3, m_3)\varphi_\epsilon^{k_1+k_2})\sigma((n_2, m_2)\varphi_\epsilon^{k_1}, (n_3, m_3)\varphi_\epsilon^{k_1+k_2}). \end{aligned}$$

□

We then have the following result.

Proposition 4.4. *The algebras $\mathbb{T}_{\Lambda, V, i} = \mathbb{T}_{\Lambda, i} \rtimes_i V$ are isomorphic to the algebras*

$$\begin{aligned}
 \mathbb{T}_{\Lambda, V, 1} &= C^*(\mathbb{Z}^2, \sigma_\theta) \rtimes_{v_1} \mathbb{Z} &= C^*(\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}, \tilde{\sigma}_\theta) \\
 &= C^*(\Lambda, \sigma_{\theta(\theta' - \theta)^{-1}}) \rtimes_{v_1} V &= C^*(S(\Lambda, V), \tilde{\sigma}_{\theta(\theta' - \theta)^{-1}}), \\
 \mathbb{T}_{\Lambda, V, 2} &= C^*(\mathbb{Z}^2, \sigma_{\theta'}) \rtimes_{v_2} \mathbb{Z} &= C^*(\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}, \tilde{\sigma}_{\theta'}) \\
 &= C^*(\Lambda, \sigma_{\theta'(\theta' - \theta)^{-1}}) \rtimes_{v_2} V &= C^*(S(\Lambda, V), \tilde{\sigma}_{\theta'(\theta' - \theta)^{-1}}).
 \end{aligned}
 \tag{4.9}$$

Proof. We just show explicitly one of the identifications. The others follow similarly. The twisted group algebra $C^*(\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}, \tilde{\sigma}_\theta)$ is generated by elements $R_{(n, m, k)}^{\tilde{\sigma}}$ satisfying

$$R_{(n, m, k)}^{\tilde{\sigma}} R_{(n', m', k')}^{\tilde{\sigma}} = \tilde{\sigma}((n, m, k), (n', m', k')) R_{(n, m, k)(n', m', k')}^{\tilde{\sigma}}.$$

The crossed product $C^*(\mathbb{Z}^2, \sigma_\theta) \rtimes_{v_1} \mathbb{Z}$ is generated by elements of the form $R_{(n, m)}^\sigma v_\epsilon^k$. The map $R_{(n, m, k)}^{\tilde{\sigma}} \mapsto R_{(n, m)}^\sigma v_\epsilon^k$ gives an identification of the generators, which also satisfies

$$\begin{aligned}
 R_{(n, m)}^\sigma v_\epsilon^k R_{(n', m')}^\sigma v_\epsilon^{k'} &= R_{(n, m)}^\sigma R_{(n', m')\varphi_\epsilon^k}^\sigma v_\epsilon^{k+k'} \\
 &= \sigma((n, m), (n', m')\varphi_\epsilon^k) R_{(n, m) + (n', m')\varphi_\epsilon^k}^\sigma v_\epsilon^{k+k'}.
 \end{aligned}$$

This gives an isomorphism $C^*(\mathbb{Z}^2, \sigma_\theta) \rtimes_{v_1} \mathbb{Z} = C^*(\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}, \tilde{\sigma}_\theta)$. \square

4.4. The magnetic Laplacian. Consider the general setting of a finitely generated discrete group Γ acting on a contractible space \tilde{X} with compact quotient $X = \tilde{X}/\Gamma$. Assume everything happens in the smooth category and we think of \tilde{X} as endowed with a metric that is invariant under the action of Γ . Upon choosing a base point $x_0 \in \tilde{X}$, we can think of the discrete set Γx_0 as a crystal of charged ions and consider the electron-ion interaction problem in \tilde{X} . This means that electrons move in \tilde{X} subject to a periodic potential. Under reasonable assumptions, one can make an *independent electron approximation* and replace the N -particle Hamiltonian with the unbounded periodic electric potential of the ion crystal with a single electron Hamiltonian in an effective periodic potential given by a bounded function (see [20] for a brief overview).

The Hamiltonian is then of the form $\Delta + V$, where the Δ is the Laplacian on \tilde{X} . We think of it as an unbounded operator on $L^2(\tilde{X})$. The Hamiltonian is invariant under translations by $\gamma \in \Gamma$, that is, $T_\gamma \Delta = \Delta T_\gamma$ and by construction V is also invariant. Here the T_γ are the operators $T_\gamma f(x) = f(x\gamma)$ on $L^2(\tilde{X})$.

One can consider on \tilde{X} a magnetic field. This is specified by a closed 2-form ω which satisfies $\gamma^* \omega = \omega$. Since \tilde{X} is contractible, there is a global magnetic potential $\omega = d\chi$. The corresponding hermitian connection $\nabla = d - i\chi$ satisfies $\nabla^2 = i\omega$. The invariance of ω implies $d(\chi - \gamma^* \chi) = 0$, so that $\chi - \gamma^* \chi = d\phi_\gamma$, where the function $\phi_\gamma(x) = \int_{x_0}^x \chi - \gamma^* \chi$ has the properties that

$$\phi_\gamma(x) - \phi_{\gamma'}(\gamma x) - \phi_{\gamma\gamma'}(x)$$

is independent of $x \in \tilde{X}$ and $\phi_\gamma(x_0) = 0$ so that

$$\sigma(\gamma, \gamma') = \exp(-i\phi_\gamma(\gamma' x_0))$$

defines a multiplier $\sigma : \Gamma \times \Gamma \rightarrow U(1)$. The Laplacian Δ is naturally replaced, in the presence of a magnetic field, by the magnetic Laplacian $\Delta^\chi = \nabla^* \nabla = (d - i\chi)^*(d - i\chi)$. This is no longer invariant under translations T_γ , but is invariant under the magnetic translations

$$(4.11) \quad T_\gamma^\phi \Delta^\chi = \Delta^\chi T_\gamma^\phi$$

where $T_\gamma^\phi f(x) = \exp(i\phi_\gamma(x))f(\gamma^{-1}x)$. Similarly, in the independent electron approximation, the effective potential V is also invariant under the magnetic translations. The magnetic translations satisfy the relations of the twisted group algebra $C_r^*(\Gamma, \bar{\sigma})$

$$T_\gamma^\phi T_{\gamma'}^\phi = \bar{\sigma}(\gamma, \gamma') T_{\gamma\gamma'}^\phi,$$

for σ as in (4.10) and $\bar{\sigma}$ the conjugate. (We refer the reader to [18], [20] for a brief overview of these well known facts.)

4.5. Discretized electron-ion interaction and Harper operators. It is usually convenient to discretize the electron-ion interaction problem. This means replacing the continuum model with Hilbert space $L^2(\tilde{X})$ by a discrete model on the Hilbert space $\ell^2(\Gamma)$. In the case without magnetic field, this is done by replacing the Laplacian Δ by its discretized version $\Delta_{discr} = r - \mathcal{R}$, where r is the cardinality of a symmetric set of generators for Γ and \mathcal{R} is the random walk operator

$$(4.12) \quad \mathcal{R} = \sum_{i=1}^r R_{\gamma_i} \quad \text{with} \quad R_{\gamma_i} f(\gamma) = f(\gamma\gamma_i)$$

for $f \in \ell^2(\Gamma)$. As in the continuum model the discretized Laplacian commutes with translations by elements $\gamma \in \Gamma$. The effective potential is then taken to be an element in the group ring $\mathbb{C}[\Gamma]$.

In the presence of a magnetic field, one can still obtain a good discretized version of the electron-ion interaction problem as in [29]. The random walk operator of (4.12) is then replaced by the Harper operator

$$(4.13) \quad \mathcal{H}_\sigma = \sum_{i=1}^r R_{\gamma_i}^\sigma,$$

with $R_{\gamma_i}^\sigma \in \mathbb{C}(\Gamma, \sigma)$ the elements of the right σ -regular representation, with σ the cocycle of (4.10). The discretized version of the magnetic Laplacian is then given by the operator

$$(4.14) \quad \Delta_{discr}^\chi = r - \mathcal{H}_\sigma,$$

which commutes with the magnetic translations $L_\gamma^{\bar{\sigma}}$. Similarly the effective potential is taken to be an element $V \in \mathbb{C}(\Gamma, \sigma)$, which then also commutes with the magnetic translations $L_\gamma^{\bar{\sigma}}$.

4.6. Harper operators for noncommutative tori and for $S(\Lambda, V)$. In the case of the noncommutative torus, viewed as the twisted group C^* -algebra $C^*(\mathbb{Z}^2, \sigma)$, the Harper operator is of the form

$$(4.15) \quad \mathcal{H}_\sigma = U + U^* + V + V^*,$$

where U and V are the generators of \mathcal{A}_θ .

The spectral theory of the Harper operator \mathcal{H}_σ of (4.15) was widely studied. In particular, it was shown in [16] that the spectrum exhibits a remarkable fractal structure (the Hofstadter butterfly) which appears to have infinitely many gaps (Cantor like spectrum) for irrational θ and finitely many gaps (band spectrum) for rational θ . The precise gap structure of the spectrum of Harper operators, as a function of the magnetic flux (that is θ in the noncommutative torus case), is a problem still under active investigation. As we see more in detail in the following, in the specific case of interest here, the gap labelling problem for the spectrum of the Harper operator is closely related to the computation of the range of the trace on the K -theory of the twisted group C^* -algebra.

In the following, we will be interested in the case of the group $S(\Lambda, V)$. In this case, after identifying it with $\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}$, the Harper operator is of the form

$$(4.16) \quad \mathcal{H}_{\tilde{\sigma}} = U + U^* + V + V^* + W + W^*,$$

where $U = R_{(0,1,0)}^{\tilde{\sigma}}$, $V = R_{(1,0,0)}^{\tilde{\sigma}}$ and $W = R_{(0,0,1)}^{\tilde{\sigma}}$.

4.7. Spectral theory and K -theory. We recall here briefly the relation between spectral theory of Harper operators and K -theory of twisted group C^* -algebras (cf. [6], [20] §3). We then proceed in the following section to analyze the specific case of $C^*(S(\Lambda, V))$.

As we have seen, the twisted group C^* -algebra $C_r^*(\Gamma, \sigma)$ is the norm closure of the twisted group ring $\mathbb{C}(\Gamma, \sigma)$ in the right σ -regular representation on $\ell^2(\Gamma)$, that is, the C^* -algebra generated by the magnetic translations R_γ^σ . If we take the weak closure of $\mathbb{C}(\Gamma, \sigma)$, we obtain the twisted group von Neumann algebra $\mathcal{U}(\Gamma, \sigma)$. Suppose given an operator $\mathcal{H}_{\sigma,V} = \mathcal{H}_\sigma + V$, with \mathcal{H}_σ the Harper operator described above and V an effective potential in $\mathbb{C}(\Gamma, \sigma)$. We have by construction $\mathcal{H}_{\sigma,V} \in \mathbb{C}(\Gamma, \sigma) \subset C_r^*(\Gamma, \sigma) \subset \mathcal{U}(\Gamma, \sigma)$, hence the spectral projections of $\mathcal{H}_{\sigma,V}$,

$$(4.17) \quad P_E = \mathbf{1}_{(-\infty, E]}(\mathcal{H}_{\sigma,V})$$

are in the von Neumann algebra, $P_E \in \mathcal{U}(\Gamma, \sigma)$. In particular, if the energy level E is not in the spectrum of $\mathcal{H}_{\sigma,V}$, then the corresponding spectral projection P_E is actually in the C^* -algebra $C_r^*(\Gamma, \sigma)$.

This implies that the question of counting gaps in the spectrum of $\mathcal{H}_{\sigma,V}$ can be reformulated as a problem of counting projections in the C^* -algebra $C_r^*(\Gamma, \sigma)$, modulo the Murray-von Neumann equivalence relation, $P \sim Q$ if there exists $V \in C_r^*(\Gamma, \sigma) \otimes \mathcal{K}$ with $P = V^*V$ and $Q = VV^*$. Equivalent spectral projections correspond to a same gap in the spectrum. The group $K_0(C_r^*(\Gamma, \sigma))$ is the Grothendieck group of the resulting abelian semi-group (with the operation of direct sum). Thus, the gap

counting problem is restated as a problem involving K -theory of C^* -algebras. More precisely, there is a faithful canonical finite trace

$$\tau(a) = \langle a\delta_1, \delta_1 \rangle_{\ell^2(\Gamma)},$$

on $C_r^*(\Gamma, \sigma)$, with δ_γ the canonical basis of $\ell^2(\Gamma)$. This extends to

$$\text{tr} = \tau \otimes \text{Tr} : \{P \in C_r^*(\Gamma, \sigma) \otimes \mathcal{K} \mid P^* = P, P^2 = P\} \rightarrow \mathbb{R},$$

with Tr the standard trace on bounded operators and induces

$$(4.18) \quad [\text{tr}] : K_0(C_r^*(\Gamma, \sigma)) \rightarrow \mathbb{R}.$$

One can obtain an estimate of the number of equivalence classes of projections by a direct computation of the range of the trace on $K_0(C_r^*(\Gamma, \sigma))$, using

$$(4.19) \quad \text{tr}(\{P \in C_r^*(\Gamma, \sigma) \otimes \mathcal{K} \mid P^* = P, P^2 = P\}) = [\text{tr}](K_0(C_r^*(\Gamma, \sigma))) \cap [0, 1].$$

5. HOMOTOPY QUOTIENT AND THE BAUM–CONNES CONJECTURE

As we show in this section, the computation of the range of the trace on K -theory is closely related to the use of the 3-manifold X_ϵ as a commutative model up to homotopy of the noncommutative space $\mathbb{T}_{\Lambda, V, i}$.

The main idea of the Baum–Connes conjecture is precisely the fact that noncommutative spaces originating from “bad quotients” have good homotopy quotients that can be used to compute geometrically invariants such as the analytic K -theory.

The group $S(\Lambda, V)$ we are considering here is a particular case of a class of groups of the form $\mathbb{Z}^2 \rtimes_\varphi \mathbb{Z}$, for some $\varphi \in \text{SL}_2(\mathbb{Z})$. The corresponding (twisted) group C^* -algebras and their K -theory were analyzed in [24]. We wish to stress here the relation between the noncommutative space and its model X_ϵ and the role of the latter in the index computations.

5.1. K -theory of $C^*(S(\Lambda, V), \tilde{\sigma})$. We now compute explicitly the K -theory of the twisted group C^* -algebra of $S(\Lambda, V)$. This can be done using the Pimsner–Voiculescu six terms exact sequence.

Lemma 5.1. *The K -theory groups of $C^*(S(\Lambda, V), \tilde{\sigma})$ are of the form*

$$(5.1) \quad \begin{aligned} K_0(C^*(S(\Lambda, V), \tilde{\sigma})) &\cong \Lambda \\ K_1(C^*(S(\Lambda, V), \tilde{\sigma})) &\cong \Lambda \oplus \Lambda / (1 - A_\epsilon)\Lambda. \end{aligned}$$

Proof. By Proposition 4.4, we can identify $C^*(S(\Lambda, V), \tilde{\sigma})$ with the crossed product $C^*(\Lambda, \sigma) \rtimes V$. Thus, we can apply the Pimsner–Voiculescu six terms exact sequence for the actions of $V \cong \mathbb{Z}$. We have

$$\begin{array}{ccccc} K_0(\mathcal{A}) & \xrightarrow{1-\alpha_*} & K_0(\mathcal{A}) & \longrightarrow & K_0(\mathcal{A} \rtimes \mathbb{Z}) \\ \uparrow \partial & & & & \downarrow \partial \\ K_1(\mathcal{A} \rtimes \mathbb{Z}) & \longleftarrow & K_1(\mathcal{A}) & \xleftarrow{1-\beta_*} & K_1(\mathcal{A}) \end{array}$$

where $\mathcal{A} = C^*(\Lambda, \sigma)$ and α_* and β_* denote, the action on $K_0(\mathcal{A})$ and $K_1(\mathcal{A})$, respectively, induced by the generator A_ϵ of the \mathbb{Z} -action on \mathcal{A} . We can identify $K_0(\mathcal{A}) = \Lambda = K_1(\mathcal{A})$. We then have $1 - \alpha_* = 0$ and $1 - \beta_* = 1 - A_\epsilon$, so that we obtain $\text{Ker}(1 - \beta_*) = \text{Ker}(1 - A_\epsilon) = 0$ and $\text{Coker}(1 - \beta_*) = \Lambda/(1 - A_\epsilon)\Lambda$. \square

We find in this way an abstract isomorphism of abelian groups

$$(5.2) \quad \begin{aligned} K_0(C^*(S(\Lambda, V)), \tilde{\sigma}) &\cong H^{odd}(X_\epsilon, \mathbb{Z}) \cong \mathbb{Z}^2 \\ K_1(C^*(S(\Lambda, V)), \tilde{\sigma}) &\cong H^{ev}(X_\epsilon, \mathbb{Z}) \cong \mathbb{Z}^2 \oplus \text{Coker}(1 - A_\epsilon). \end{aligned}$$

This identification can be justified more naturally in terms of the Baum–Connes conjecture, as we discuss in the following.

5.2. K -theory and the twist. The following result shows that the presence of the twisting by $\tilde{\sigma}$ has no effect on the K -theory.

Lemma 5.2. *There is an isomorphism*

$$(5.3) \quad K_i(C^*(S(\Lambda, V), \tilde{\sigma})) \cong K_i(C^*(S(\Lambda, V)))$$

between the K -theory of the twisted group C^ -algebra $C^*(S(\Lambda, V), \tilde{\sigma})$ and the K -theory of the untwisted $C^*(S(\Lambda, V))$.*

Proof. The argument is similar to that used in [18], [19] and Corollary 2.2 of [13]. The cocycle $\tilde{\sigma}$ is real in the sense of Definition 1.12 of [13], being of the form (4.8), with σ of the exponential form $\sigma((n, m), (n', m')) = \exp(-\pi i \theta(mn' - nm'))$. Thus, as in Corollary 1.13 of [13], the identification (5.3) follows using a homotopy $\exp(-\pi i \theta(mn' - nm'))$, with $t \in [0, 1]$. \square

Notice in fact that for groups of the form $\Gamma = \mathbb{Z}^2 \rtimes_\varphi \mathbb{Z}$, with $\varphi \in \text{SL}_2(\mathbb{Z})$ all cocycles $\sigma : \Gamma \times \Gamma \rightarrow U(1)$ are real in the above sense. This was observed already in [24].

Lemma 5.3. *Let $\sigma \in Z^2(\Gamma, U(1))$ be a cocycle. Then σ is cohomologous to a real cocycle, that is, to an element of $Z^2(\Gamma, U(1))$ that is of the form $\exp(2\pi i \zeta)$ for $\zeta \in Z^2(\Gamma, \mathbb{R})$.*

Proof. We can see it easily as in §2.2 of [18], by considering the exact sequence of coefficient groups

$$1 \rightarrow \mathbb{Z} \xrightarrow{\iota} \mathbb{R} \xrightarrow{\exp(2\pi i \cdot)} U(1) \rightarrow 1$$

and the long exact cohomology sequence

$$(5.4) \quad \cdots \rightarrow H^2(\Gamma, \mathbb{Z}) \rightarrow H^2(\Gamma, \mathbb{R}) \xrightarrow{\exp(2\pi i \cdot)^*} H^2(\Gamma, U(1)) \xrightarrow{\delta} H^3(\Gamma, \mathbb{Z}) \xrightarrow{\iota_*} H^3(\Gamma, \mathbb{R}) \rightarrow \cdots$$

Since in our case, for $\Gamma = S(\Lambda, V)$, we have $E\Gamma = \mathbb{R}^2 \rtimes \mathbb{R}$ and $B\Gamma = X_\epsilon$, we see that

$$H^2(\Gamma, \mathbb{Z}) = H^2(X_\epsilon, \mathbb{Z}) = H_1(X_\epsilon, \mathbb{Z}) = \mathbb{Z} \oplus \Lambda/(1 - A)\Lambda$$

and

$$H^3(\Gamma, \mathbb{Z}) = H^3(X_\epsilon) = \mathbb{Z}.$$

We then see that, in the sequence (5.4) the map ι_* is injective so that $\delta = 0$. Thus, all elements in $H^2(\Gamma, U(1))$ come from $H^2(\Gamma, \mathbb{R})$ via the exponential map. \square

In fact, we do not need this general fact, as the cocycle we are using is already constructed in the desired exponential form, but we stated it here for completeness.

5.3. Thom isomorphism, homotopy quotients and Baum–Connes. It is known that the group $S(\Lambda, V)$ satisfies the Baum–Connes conjecture (with coefficients). In fact the group $\mathrm{SL}_2(\mathbb{Z})$ is known to satisfy the Baum–Connes conjecture with coefficients, hence by [8] so does the group $\mathbb{Z}^2 \rtimes_{\varphi_\epsilon} \mathbb{Z}$ with $\varphi_\epsilon \in \mathrm{SL}_2(\mathbb{Z})$.

This means that the Kasparov assembly map is an isomorphism, hence the K -theory of the C^* -algebra $C^*(S(\Lambda, V))$ can be computed in terms of the geometric K -theory of the homotopy quotient $\underline{B}\Gamma$, the classifying space for proper actions (cf. [4]). This relates directly the analytic K -theory of the C^* -algebra to the topological K -theory of the 3-manifold X_ϵ .

Lemma 5.4. *The Kasparov assembly map for $C^*(S(\Lambda, V))$ gives an isomorphism*

$$(5.5) \quad \begin{aligned} \mu : K^1(X_\epsilon) &\xrightarrow{\cong} K_0(C^*(S(\Lambda, V))) \\ \mu : K^0(X_\epsilon) &\xrightarrow{\cong} K_1(C^*(S(\Lambda, V))). \end{aligned}$$

Proof. In our case the space $\underline{E}\Gamma$ is the solvable Lie group $S(\mathbb{R}^2, \mathbb{R}) = \mathbb{R}^2 \rtimes \mathbb{R}$ and the homotopy quotient $\Gamma \backslash \underline{B}\Gamma$ is the 3-manifold $X_\epsilon = S(\Lambda, V) \backslash S(\mathbb{R}^2, \mathbb{R})$. This can be identified with the mapping torus

$$X_\epsilon = T^2 \times [0, 1] / ((x, y), 0) \sim (A_\epsilon(x, y), 1).$$

For a mapping torus, the Thom isomorphism [10] gives the identification

$$(5.6) \quad K_{i+1}(C(X_\epsilon)) = K_i(C(T^2) \rtimes_{A_\epsilon} \mathbb{Z}).$$

Moreover, the C^* -algebra $C(T^2) \rtimes_{A_\epsilon} \mathbb{Z}$ is identified with $C^*(\Lambda) \rtimes V$ by Fourier transform, which identifies $C(T^2) = C^*(\Lambda)$ for $T^2 = \mathbb{R}^2/\Lambda$. The algebra $C^*(\Lambda) \rtimes V$ is then isomorphic to $C^*(S(\Lambda, V))$, by the same argument of Proposition 4.4 in the untwisted case. \square

6. TWISTED INDEX THEOREM, K -THEORY, AND THE RANGE OF THE TRACE

As we have seen, the 2-cocycle $\tilde{\sigma}$ on $\Gamma = S(\Lambda, V)$, is of the form $\tilde{\sigma} = \exp(2\pi i \zeta)$, with $\zeta \in H^2(\Gamma, \mathbb{R})$. Upon identifying $H^2(\Gamma, \mathbb{R}) = H^2(B\Gamma, \mathbb{R}) = H^2(X_\epsilon, \mathbb{R})$, we can identify the 2-cocycle ζ with a closed 2-form ω_ϵ on the 3-manifold X_ϵ . We denote by $\tilde{\omega}_\epsilon$ its pullback to the universal covering $\tilde{X}_\epsilon = S(\mathbb{R}^2, \mathbb{R})$. This is a Γ -invariant 2-form, $\gamma^* \omega_\epsilon = \omega_\epsilon$, which we previously interpreted as a magnetic field.

Lemma 6.1. *The real 2-cocycle $\zeta \in H^2(\Gamma, \mathbb{R})$ with $\tilde{\sigma} = \exp(2\pi i \zeta)$ is given by*

$$(6.1) \quad \zeta((\lambda, k), (\eta, r)) = \frac{1}{4\pi i} \int_{\mathcal{R}} \omega,$$

where ω is the closed 2-form on $T^2 = \mathbb{R}^2/\Lambda$ associated to the cocycle σ on Λ , with magnetic flux $\int_{T^2} \omega = 2\pi i \theta(\theta' - \theta)^{-1}$, and $\mathcal{R} \subset \mathbb{R}^2$ is the oriented parallelogram with vertices

$$(6.2) \quad \{0, A_\epsilon^k(\eta), \lambda, \lambda + A_\epsilon^k(\eta)\}.$$

Proof. On \tilde{X}_ϵ the form $\tilde{\omega}_\epsilon$ is exact, hence we have a global magnetic potential χ_ϵ with $\tilde{\omega}_\epsilon = d\chi_\epsilon$ and $d(\chi_\epsilon - \gamma^*\chi_\epsilon) = 0$, or $\chi_\epsilon - \gamma^*\chi_\epsilon = d\phi_\gamma$, as before, where the ϕ_γ recovers the cocycle $\tilde{\sigma}$ by the formula

$$\tilde{\sigma}(\gamma, \gamma') = \exp(-\phi_\gamma(\gamma'x_0)) = \exp\left(\int_{x_0}^{\gamma'x_0} \gamma^*\chi_\epsilon - \chi_\epsilon\right).$$

We know from Lemma 4.3 that the cocycle $\tilde{\sigma}$ has the form $\tilde{\sigma}_u((\lambda, k), (\eta, r)) = \sigma_u(\lambda, A_\epsilon^k(\eta))$, for $u = \theta(\theta' - \theta)^{-1}$, so that we have

$$\zeta((\lambda, k), (\eta, r)) = \frac{\theta(\theta' - \theta)^{-1}}{2} A_\epsilon^k(\eta) \wedge \lambda,$$

that is, $\zeta((\lambda, k), (\eta, r)) = \xi(\lambda, A_\epsilon^k(\eta))$, where

$$(6.3) \quad \xi(\lambda, A_\epsilon^k(\eta)) = \frac{1}{2\pi i} \int_0^{A_\epsilon^k(\eta)} U_\lambda^* \chi - \chi.$$

Here χ is the magnetic potential on \mathbb{R}^2 associated to the closed 2-form ω with

$$\int_{T^2} \omega = 2\pi i \theta(\theta' - \theta)^{-1}.$$

Let then \mathcal{R} denote the oriented parallelogram in \mathbb{R}^2 with vertices as in (6.2). We have

$$(6.4) \quad \frac{1}{2\pi i} \int_{\mathcal{R}} \omega = \frac{1}{2\pi i} \int_0^{A_\epsilon^k(\eta)} (U_\lambda^* \chi - \chi) - \frac{1}{2\pi i} \int_0^\lambda (U_{A_\epsilon^k(\eta)}^* \chi - \chi).$$

Using the fact that $\xi(\eta, \lambda) = -\xi(\lambda, \eta)$, this gives

$$\frac{1}{2\pi i} \int_{\mathcal{R}} \omega = 2\zeta((\lambda, k), (\eta, r)).$$

□

6.1. Spectral flow and odd Chern character. An element of $K_1(C(X_\epsilon))$ can be viewed as the class $[g]$ of $g \in U_N(C(X_\epsilon))$, which we can see as a differentiable map $g : X_\epsilon \rightarrow \text{GL}_N(\mathbb{C})$. We proceed as in [14] and we consider the associated 1-form

$$(6.5) \quad \beta(g) = g^{-1}dg \in \Omega^1(X_\epsilon, gl_N(\mathbb{C})).$$

The corresponding family of connections $\nabla_u = d + u\beta(g)$ on the trivial bundle $X_\epsilon \times \mathbb{C}^N$ determines a closed Chern-Simons form

$$(6.6) \quad Ch(g) := cs(d, d + \beta(g)) = \int_0^1 \text{Tr}\left(\frac{d}{du}(\nabla_u) e^{\nabla_u^2}\right) du,$$

which gives the odd Chern character $Ch(g)$. As shown in [14], this has an expression as an odd differential form

$$(6.7) \quad Ch(g) = \sum_{k=0}^{\infty} (-1)^k \frac{k!}{(2k+1)!} \text{Tr}(\beta(g)^{2k+1}).$$

One then has, see [14], that the pairing

$$(6.8) \quad \langle D, [g] \rangle = SF(D, g^{-1}Dg)$$

of an odd Fredholm module (\mathcal{H}, D) with $[g] \in K_1$ is given by the spectral flow along $D_u = (1 - u)D + ug^{-1}Dg$. In the case where $D = \not{D}$ is the Dirac operator of a compact spin manifold, this is computed by the Atiyah–Patodi–Singer index formula [3]. In our case, this gives

$$(6.9) \quad SF(\not{D}, g^{-1}\not{D}g) = -\frac{1}{(2\pi i)^2} \int_{X_\epsilon} \hat{A}(X_\epsilon) Ch(g).$$

6.2. Twisted index theorem. We need the twisted version of (6.9) above. Let $\not{D} = \not{D}_{X_\epsilon}$ be the Dirac operator on X_ϵ , and let $\tilde{\not{D}}$ be its lift to the universal cover $\tilde{X}_\epsilon = S(\mathbb{R}^2, \mathbb{R})$. We then consider the twisting $\tilde{\not{D}} \otimes \nabla$ of the operator $\tilde{\not{D}}$ by the hermitian connection $\nabla = d + i\eta_\epsilon$ on the trivial line bundle on \tilde{X}_ϵ , with η_ϵ the 1-form giving the magnetic potential $d\eta_\epsilon = \omega_\epsilon$ on \tilde{X}_ϵ .

While the operator $\tilde{\not{D}}$ is Γ -invariant, with $\Gamma = S(\Lambda, V)$, the twisted operator $\tilde{\not{D}} \otimes \nabla$ is only invariant under the projective action $(\Gamma, \tilde{\sigma})$ of the magnetic translations $R_\gamma^{\tilde{\sigma}}$. Consider then the 1-parameter family of operators $D_u = \tilde{\not{D}}_u \otimes \nabla$, where $\tilde{\not{D}}_u = (1 - u)\tilde{\not{D}} + ug^{-1}\tilde{\not{D}}g$, for $[g] \in K^1(X_\epsilon)$ and the associated operator $\mathcal{D}_g = \frac{\partial}{\partial u} + D_u$ on $\tilde{X}_\epsilon \times [0, 1]$, which we can extend to $\tilde{X}_\epsilon \times \mathbb{R}$ (cf. p.95 of [3]).

Theorem 6.2. *The range of the trace on $K_0(C^*(S(\Lambda, V), \tilde{\sigma}))$ is given by*

$$(6.10) \quad [\text{tr}](\mu_{\tilde{\sigma}}[g]) = \frac{-1}{(2\pi i)^2} \int_{X_\epsilon} \hat{A} e^{\omega_\epsilon} Ch(g),$$

where $\mu_{\tilde{\sigma}} : K^1(X_\epsilon) \rightarrow K_0(C^*(S(\Lambda, V), \tilde{\sigma}))$ is the (twisted) Kasparov isomorphism, $[g] \in K^1(X_\epsilon)$, and ω_ϵ is the closed 2-form on X_ϵ associated to the cocycle $\tilde{\sigma}$.

Proof. We let P^\pm be the projections on the L^2 -kernel of $\mathcal{D}_g \mathcal{D}_g^*$ and $\mathcal{D}_g^* \mathcal{D}_g$, respectively, namely

$$\mathcal{D}_g P^+ = 0 \quad \mathcal{D}_g^* P^- = 0.$$

The P^\pm have smooth kernels $P^\pm(x, y)$ and the $(\Gamma, \tilde{\sigma})$ -invariance of $\tilde{\not{D}} \otimes \nabla$ implies that

$$e^{-i\phi_\gamma(x)} P^\pm(\gamma x, \gamma y) e^{i\phi_\gamma(y)} = P^\pm(x, y),$$

which implies that $P^\pm(x, x)$ is Γ -invariant, for $\Gamma = S(\Lambda, V)$.

We proceed as in [1] and consider the von Neumann trace

$$\text{tr}(P^\pm) = \int_{X_\epsilon \times S^1} \text{tr} P^\pm((x, t), (x, t)) dx dt,$$

where $\text{tr} P^\pm(x, x)$ is the pointwise trace. The L^2 -index of \mathcal{D}_g is given by

$$(6.11) \quad \text{Ind}_{L^2}(\mathcal{D}_g) = \text{tr}(P^+) - \text{tr}(P^-).$$

We define \bar{P}^\pm by the smooth kernels

$$(6.12) \quad \bar{P}^\pm(x, y) = \int_{S^1} \text{tr} P^\pm((x, t), (y, t)) dt.$$

These satisfy $\text{tr}(\bar{P}^\pm) = \text{tr}(P^\pm)$ by

$$\int_{X_\epsilon \times S^1} \text{tr} P^\pm((x, t), (x, t)) dx dt = \int_{X_\epsilon} \text{tr} \bar{P}^\pm(x, x) dx.$$

The projections \bar{P}^\pm are in the von Neumann algebra $\mathcal{U}(\Gamma, \tilde{\sigma})$. After adding a compact perturbation in $C_r^*(\Gamma, \tilde{\sigma})$ one obtains a well defined index (cf. [22], [27]),

$$(6.13) \quad \text{Ind}_{(\Gamma, \tilde{\sigma})}(\mathcal{D}_g) = [\bar{P}^+] - [\bar{P}^-] \in K_0(C_r^*(\Gamma, \tilde{\sigma})).$$

The (twisted) Kasparov map $\mu : K^1(X_\epsilon) \rightarrow K_0(C^*(\Gamma, \tilde{\sigma}))$ is given by

$$(6.14) \quad \mu_{\tilde{\sigma}}[g] = \text{Ind}_{(\Gamma, \tilde{\sigma})}(\mathcal{D}_g).$$

We obtain in this way that

$$\text{Ind}_{L^2}(\mathcal{D}_g) = \text{tr}(\bar{P}^+) - \text{tr}(\bar{P}^-) = \text{tr}(\text{Ind}_{(\Gamma, \tilde{\sigma})}(\mathcal{D}_g)).$$

Consider the heat kernel $e^{-t\mathcal{D}^2}$, where

$$\mathcal{D} = \begin{pmatrix} 0 & \mathcal{D}_g^* \\ \mathcal{D}_g & 0 \end{pmatrix} \quad \text{with} \quad \mathcal{D}^2 = \begin{pmatrix} \mathcal{D}_g^* \mathcal{D}_g & 0 \\ 0 & \mathcal{D}_g \mathcal{D}_g^* \end{pmatrix}.$$

We have

$$\lim_{t \rightarrow \infty} \text{tr}_s(e^{-t\mathcal{D}^2}) = \text{tr}(P^+) - \text{tr}(P^-)$$

and

$$\frac{\partial}{\partial t} \text{tr}_s(e^{-t\mathcal{D}^2}) = -\text{tr}_s(\mathcal{D}^2 e^{-t\mathcal{D}^2}) = \text{tr}_s([\mathcal{D} e^{-t\mathcal{D}^2}, \mathcal{D}]) = 0.$$

Thus

$$\begin{aligned} \text{tr}(P^+) - \text{tr}(P^-) &= \lim_{t \rightarrow \infty} \text{tr}_s(e^{-t\mathcal{D}^2}) = \lim_{t \rightarrow 0} \text{tr}_s(e^{-t\mathcal{D}^2}) \\ &= \frac{-1}{(2\pi i)^2} \int_{X_\epsilon \times S^1} \hat{A} Ch(\nabla_u) = \frac{-1}{(2\pi i)^2} \int_{X_\epsilon} \hat{A} e^{\omega_\epsilon} Ch(g), \end{aligned}$$

where $Ch(\nabla_u) = \text{tr}(\beta e^{(d+u\beta)^2})$ for $\beta = g^{-1}dg$, with $\int_{S^1} Ch(\nabla_u) = Ch(g)$. \square

6.3. Range of the trace. Using the twisted index theorem we can then compute explicitly the range of the trace on $K_0(C^*(S(\Lambda, V), \tilde{\sigma}))$. We obtain the following result.

Proposition 6.3. *The range of the trace on $K_0(C^*(S(\Lambda, V), \tilde{\sigma}))$ is*

$$(6.15) \quad [\text{tr}](K_0(C^*(S(\Lambda, V), \tilde{\sigma}))) = \mathbb{Z} + \mathbb{Z}\theta(\theta' - \theta)^{-1}.$$

Proof. Since X_ϵ is a 3-manifold, when we expand the terms in the cohomological formula (6.10) as

$$\begin{aligned} \hat{A}(X_\epsilon) &= 1 - \frac{1}{24} p_1(X_\epsilon) + \cdots \\ e^{\omega_\epsilon} &= 1 + \omega_\epsilon + \frac{1}{2} \omega_\epsilon^2 + \cdots \\ Ch(g) &= -\frac{1}{6} \text{Tr}(\beta(g)) + \frac{1}{5!} \text{Tr}(\beta^3(g)) + \cdots, \end{aligned}$$

only the terms of the wedge product $\hat{A}(X_\epsilon) e^\omega Ch(g)$ that give differential forms of order up to 3 can contribute nontrivially.

Thus, we obtain the terms

$$(6.16) \quad \frac{1}{(2\pi)^2} \int_{X_\epsilon} \left(\frac{-1}{6} \text{Tr}(\beta(g)) \wedge \omega + \frac{1}{5!} \text{Tr}(\beta(g)^3) \right).$$

The term

$$\frac{1}{(2\pi)^2} \int_{X_\epsilon} \frac{1}{5!} \text{Tr}(\beta(g)^3) = \frac{1}{(2\pi)^2} \int_{X_\epsilon} Ch(g)$$

is the term one would find in the untwisted case, and it gives the untwisted odd Chern character.

For the remaining term

$$\frac{1}{(2\pi)^2} \frac{-1}{6} \int_{X_\epsilon} \text{Tr}(\beta(g)) \wedge \omega_\epsilon$$

the range as $[g]$ varies in $K_1(C(X_\epsilon))$ is given by $\mathbb{Z}R(\omega)$, where $R(\omega)$ is the range of the linear form

$$T_\omega : [g] \mapsto \frac{1}{(2\pi)^2} \frac{-1}{6} \int_{X_\epsilon} \text{Tr}(\beta(g)) \wedge \omega_\epsilon \in \mathbb{R}.$$

First notice that, with the notation $Ch_1(g) = \frac{-1}{6} \text{Tr}(\beta(g))$, we have

$$\int_C Ch_1(g) = 2\pi i \deg(g|_C) \in 2\pi i \mathbb{Z},$$

for $C \in H_1(X_\epsilon, \mathbb{Z})$. Thus, we obtain

$$\frac{1}{2\pi i} \int_{X_\epsilon} Ch_1(g) \wedge PD(C) \in \mathbb{Z},$$

for $PD(C) \in H^2(X_\epsilon, \mathbb{Z}) \hookrightarrow H^2(X_\epsilon, \mathbb{R})$. Now consider the explicit description of the 2-form ω_ϵ given in Lemma 6.1 above. We can write

$$\omega_\epsilon = 2\pi i \theta (\theta' - \theta)^{-1} \bar{\omega}_\epsilon,$$

where $\bar{\omega}_\epsilon \in H^2(X_\epsilon, \mathbb{Z})$ is given by

$$\bar{\omega}_\epsilon(v, w) = A_\epsilon^k(\eta) \wedge \lambda,$$

for $v = ((0, 0), (\lambda, k))$ and $w = ((0, 0), (\eta, r))$. Thus, we see that we can write

$$\begin{aligned} \frac{1}{(2\pi)^2} \int_{X_\epsilon} Ch_1(g) \wedge \omega_\epsilon &= -\frac{1}{2\pi i} \theta (\theta' - \theta)^{-1} \int_{X_\epsilon} Ch_1(g) \wedge \bar{\omega}_\epsilon \\ &= -\frac{1}{2\pi i} \theta (\theta' - \theta)^{-1} \int_{PD(\bar{\omega}_\epsilon)} Ch_1(g) = \theta (\theta' - \theta)^{-1} \deg(g|_{PD(\bar{\omega}_\epsilon)}) \in \theta (\theta' - \theta)^{-1} \mathbb{Z}. \end{aligned}$$

□

7. ISOSPECTRAL DEFORMATIONS AND SPECTRAL TRIPLES

In noncommutative geometry, the analog of Riemannian structures is provided by the formalism of spectral triples [11]. A spectral triple on a noncommutative space \mathcal{A} (where \mathcal{A} is a C^* -algebra) consists of the data $(\mathcal{A}_\infty, \mathcal{H}, D)$ of a dense involutive subalgebra \mathcal{A}_∞ , a representation $\pi : \mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ as bounded operators on a Hilbert space \mathcal{H} and a self-adjoint operator D on \mathcal{H} , with compact resolvent, satisfying the compatibility condition

$$(7.1) \quad [D, \pi(a)] \in \mathcal{B}(\mathcal{H}), \quad \forall a \in \mathcal{A}_\infty.$$

In particular, in the commutative case, to a Riemannian *spin*-manifold X one can associate a canonical spectral triple $(C^\infty(X), L^2(X, S), \not{D})$. A reconstruction theorem [25] shows that a spectral triple where the algebra is abelian, which satisfies a list of axioms, is the canonical spectral triple of a Riemannian *spin*-manifold.

In our case, we have a spectral triple associated to the 3-manifold X_ϵ , where the spinor bundle is a complex 2-plane bundle and the Dirac operator can be written in the form

$$(7.2) \quad \not{D}_{X_\epsilon} = c(dt) \frac{\partial}{\partial t} + c(e^t dx) \frac{\partial}{\partial x} + c(e^{-t} dy) \frac{\partial}{\partial y},$$

where $\{dt, e^t dx, e^{-t} dy\}$ is the basis of the cotangent bundle of $S(\mathbb{R}^2, \mathbb{R}, \epsilon) = \mathbb{R}^2 \rtimes_\epsilon \mathbb{R}$ and $c(\omega)$ denotes Clifford multiplication by the 1-form ω .

More explicitly, (7.2) is of the form

$$(7.3) \quad \not{D}_{X_\epsilon} = \frac{\partial}{\partial t} \sigma_0 + e^t \frac{\partial}{\partial x} \sigma_1 + e^{-t} \frac{\partial}{\partial y} \sigma_2 = \begin{pmatrix} \frac{\partial}{\partial t} & e^{-t} \frac{\partial}{\partial y} - ie^t \frac{\partial}{\partial x} \\ e^{-t} \frac{\partial}{\partial y} + ie^t \frac{\partial}{\partial x} & -\frac{\partial}{\partial t} \end{pmatrix},$$

where σ_i , for $i = 0, 1, 2$, are the Pauli matrices.

Our purpose here is to show that this commutative spectral triple can be deformed isospectrally to a spectral triple for the noncommutative tori $\mathbb{T}_{\Lambda, i}$.

7.1. The Connes–Landi isospectral deformations. We consider the problem from the point of view of the Connes–Landi isospectral deformations [12]. This provides a general procedure to deform commutative spectral triples to noncommutative ones isospectrally, for manifolds with isometric torus actions.

We recall briefly the construction of isospectral deformations, in a version that is best adapted to our setting.

Suppose given a spectral triple $(C^\infty(X), L^2(X, S), \not{D}_X)$ associated to a compact Riemannian *spin*-manifold X . Assume that the manifold X has an action of a torus T^2 by isometries, $T^2 \subset \text{Isom}(X)$. Then one considers a noncommutative algebra \mathcal{A}_θ , depending on a real parameter $\theta \in \mathbb{R}$, which is obtained by decomposing the operators $\pi(f) \in \mathcal{B}(\mathcal{H})$, for $f \in C^\infty(X)$ and $\mathcal{H} = L^2(X, S)$ according to their weighted components

$$(7.4) \quad \pi(f) = \sum_{n, m \in \mathbb{Z}} \pi(f_{n, m}),$$

where

$$(7.5) \quad \alpha_\tau(\pi(f_{n,m})) = e^{2\pi i(n\tau_1 + m\tau_2)} \pi(f_{n,m}), \quad \forall \tau = (\tau_1, \tau_2) \in T^2,$$

for

$$(7.6) \quad \alpha_\tau(T) = U(\tau)TU(\tau)^*, \quad \forall T \in \mathcal{B}(\mathcal{H}), \forall \tau \in T^2,$$

with $U(\tau)$ the unitary transformations implementing the T^2 -action on $\mathcal{H} = L^2(X, S)$ by

$$U(\tau)\psi(x) = \psi(\tau^{-1}(x)).$$

Let L_1 and L_2 denote the infinitesimal generators of the action

$$(7.7) \quad U(\tau) = \exp(2\pi i\tau L) = \exp(2\pi i(\tau_1 L_1 + \tau_2 L_2)).$$

We consider then the subalgebra of $\mathcal{B}(\mathcal{H})$ generated by the operators of the form

$$(7.8) \quad \pi_{\xi_1, \xi_2}(f) = \sum_{n,m} \pi(f_{n,m}) e^{-2\pi i(\xi_1 n L_2 + \xi_2 m L_1)},$$

where ξ_1 and ξ_2 are two real parameters.

Lemma 7.1. *For homogeneous operators $\pi(f)_{n,m}$ define the deformed product*

$$(7.9) \quad f_{n,m} *_{\xi_1, \xi_2} h_{k,r} := e^{-2\pi i(\xi_1 n r + \xi_2 m k)} f_{n,m} h_{k,r}.$$

The product of operators of the form (7.8) satisfies

$$\pi_{\xi_1, \xi_2}(f_{n,m}) \pi_{\xi_1, \xi_2}(h_{k,r}) = \pi_{\xi_1, \xi_2}(f_{n,m} *_{\xi_1, \xi_2} h_{k,r}).$$

Proof. One checks directly that the operator product $\pi(f_{\xi_1, \xi_2})\pi(h_{\xi_1, \xi_2})$ is given in components by

$$\pi(f_{n,m}) *_{\xi_1, \xi_2} \pi(h_{k,r}) = e^{-2\pi i(\xi_1 n r + \xi_2 m k)} \pi(f_{n,m}) \pi(h_{k,r}).$$

□

One can recognize in (7.9) the convolution product of the twisted group C^* -algebra $C^*(\mathbb{Z}^2, \sigma)$ with the cocycle

$$\sigma((n, m), (k, r)) = \exp(-2\pi i(\xi_1 n r + \xi_2 m k)).$$

As shown in [12], the operators (7.8) have bounded commutators with the Dirac operator. In fact, since T^2 acts by isometries, the Dirac operator satisfies

$$U(\tau)DU(\tau)^* = D,$$

i.e. it is of bidegree $(0, 0)$. Thus, one sees that the commutators

$$\begin{aligned} [D, \pi_{\xi_1, \xi_2}(f)] &= \sum_{n,m} [D, \pi(f)_{n,m} e^{-2\pi i(\xi_1 n L_2 + \xi_2 m L_1)}] \\ &= \sum_{n,m} [D, \pi(f)]_{n,m} e^{-2\pi i(\xi_1 n L_2 + \xi_2 m L_1)}, \end{aligned}$$

which is still a bounded operator on \mathcal{H} .

We consider in particular the case where $\xi_2 = u/2 = -\xi_1$. We denote by $\mathcal{A}_u = C^\infty(X)_u$ the deformed algebra, that is, the algebra generated by the (7.8). The deformed spectral triple is given by the data $(\mathcal{A}_u, L^2(X, S), \not{D}_X)$.

7.2. Noncommutative solvmanifolds. We apply the procedure described above to obtain an isospectral deformation of the solvmanifold X_ϵ , which corresponds to deforming the fiber tori to noncommutative tori.

The canonical spectral triple for X_ϵ consists of the data $(C^\infty(X_\epsilon), L^2(X_\epsilon, S), \not{D}_{X_\epsilon})$, with the Dirac operator of the form (7.2).

There is a torus action on X_ϵ by isometries, which consists of translations along the fibers of the fibration $T^2 \rightarrow X_\epsilon \rightarrow S^1$. This acts on spinors by unitaries

$$(7.10) \quad U(\tau)\psi((x, y), t) = \psi((x + e^t\tau_1, y + e^{-t}\tau_2), t),$$

for $\tau \in T^2 = \mathbb{R}^2/\Lambda$ and $(x, y) \in T_t^2 = \mathbb{R}^2/\Lambda_t$, the fiber over $t \in S^1$, with $(e^t\lambda_1, e^{-t}\lambda_2) \in \Lambda_t$, for $(\lambda_1, \lambda_2) \in \Lambda$.

The action clearly preserves the metric $dt^2 + e^t dx^2 + e^{-t} dy^2$ hence the Dirac operator (7.3) satisfies

$$U(\tau)\not{D}_{X_\epsilon}U(\tau)^* = \not{D}_{X_\epsilon}.$$

The infinitesimal generators of the action α_τ are the operators $2\pi L_1 = e^t \frac{\partial}{\partial x}$, $2\pi L_2 = e^{-t} \frac{\partial}{\partial y}$ with $U(\tau) = \exp(2\pi i(\tau_1 L_1 + \tau_2 L_2))$.

We introduce the following notation. We denote by E_λ , for $\lambda \in \Lambda$, the function

$$(7.11) \quad E_\lambda((x, y), t) := e^{2\pi i \langle \Theta_{-t}(x, y), \lambda \rangle},$$

where, as above, $\Theta_{-t}(x, y) = (e^{-t}x, e^t y)$ and $\langle (a, b), \lambda \rangle = a\lambda_1 + b\lambda_2$. We also denote by $\Xi_u(\lambda, L_1, L_2)$ the operator

$$(7.12) \quad \Xi_u(\lambda, L_1, L_2) := \exp\left(i\pi \frac{u}{(\theta' - \theta)} \lambda \wedge (L_1, L_2)\right),$$

acting on $\mathcal{H} = L^2(X_\epsilon, S)$.

Proposition 7.2. *The deformed algebra $C^\infty(X_\epsilon)_u$, for $u \in \mathbb{R}$, is the C^* -subalgebra of $\mathcal{B}(\mathcal{H})$, with $\mathcal{H} = L^2(X_\epsilon, S)$ generated by the operators of the form*

$$(7.13) \quad \pi_u(f) = E_\lambda \Xi_u(\lambda, L_1, L_2).$$

Proof. The induced action $\alpha : T^2 \rightarrow \text{Aut}(C^\infty(X_\epsilon))$ defined by

$$\pi(\alpha_\tau(f)) = U(\tau)\pi(f)U(\tau)^*$$

is of the form $\alpha_\tau(f)((x, y), t) = f((x + e^t\tau_1, y + e^{-t}\tau_2), t)$.

Thus, a homogeneous operator of bidegree $\lambda = (\lambda_1, \lambda_2)$ is in this case a function $f_\lambda((x, y), t)$ with the property that

$$(7.14) \quad \alpha_\tau(f_\lambda)((x, y), t) = e^{2\pi i(\lambda_1\tau_1 + \lambda_2\tau_2)} f_\lambda((x, y), t).$$

This condition is satisfied by functions of the form

$$(7.15) \quad f_\lambda((x, y), t) = \exp(2\pi i \langle \Theta_{-t}(x, y), \lambda \rangle) = \exp(2\pi i(e^{-t}\lambda_1 x + e^t\lambda_2 y)).$$

Under the change of variables

$$(7.16) \quad \mathbb{Z}^2 \rightarrow \Lambda, \quad (n, m) \mapsto \lambda = (n + m\theta, n + m\theta'),$$

the condition (7.14) corresponds to elements $f_{n,m}$ of bidegree (n, m) for the corresponding action of $T^2 = \mathbb{R}^2/\mathbb{Z}^2$. Thus, using this change of coordinates to pass in

(7.8) from \mathbb{Z}^2 to Λ , we can see that elements of the deformed algebra of the form (7.8) correspond to elements of the form

$$\sum_{\lambda} a_{\lambda} E_{\lambda} \Xi_u(\lambda, L_1, L_2),$$

for $\xi_2 = u/2 = -\xi_1$. □

Set $2u_{\theta} = u/(\theta' - \theta)$. The operators (7.13) act on spinors by

$$\begin{aligned} (\pi_u(f)\psi)((x, y), t) &= E_{\lambda}((x, y), t) (U((- \lambda_2 u_{\theta}, \lambda_1 u_{\theta}))\psi)((x, y), t) \\ &= e^{2\pi i \langle \Theta_{-t}(x, y), \lambda \rangle} \psi((x - e^t \lambda_2 u_{\theta}, y + e^{-t} \lambda_1 u_{\theta}), t). \end{aligned}$$

Proposition 7.3. *The operators*

$$(7.17) \quad \pi(R_{\lambda}^{\sigma}) := E_{\lambda} \Xi_u(\lambda, L_1, L_2)$$

define a representation on $\mathcal{H} = L^2(X_{\epsilon}, S)$ of the noncommutative torus $C^*(\Lambda, \sigma)$, with the cocycle

$$\sigma(\lambda, \eta) = \exp(2\pi i u_{\theta} \lambda \wedge \eta).$$

Proof. Notice that we have

$$U(-\lambda_2 u_{\theta}, \lambda_1 u_{\theta}) e^{2\pi i \langle \Theta_{-t}(x, y), \eta \rangle} = e^{2\pi i u_{\theta} \lambda \wedge \eta} e^{2\pi i \langle \Theta_{-t}(x, y), \eta \rangle}.$$

Thus, we obtain

$$\begin{aligned} e^{2\pi i \langle \Theta_{-t}(x, y), \lambda \rangle} \Xi_u(\lambda, L_1, L_2) e^{2\pi i \langle \Theta_{-t}(x, y), \eta \rangle} \Xi_u(\eta, L_1, L_2) &= \\ e^{2\pi i u_{\theta} \lambda \wedge \eta} e^{2\pi i \langle \Theta_{-t}(x, y), \lambda \rangle} e^{2\pi i \langle \Theta_{-t}(x, y), \eta \rangle} \Xi_u(\lambda, L_1, L_2) \Xi_u(\eta, L_1, L_2) &= \\ e^{2\pi i u_{\theta} \lambda \wedge \eta} e^{2\pi i \langle \Theta_{-t}(x, y), \lambda + \eta \rangle} \Xi_u(\lambda + \eta, L_1, L_2). \end{aligned}$$

This shows that the operators $\pi(R_{\lambda}^{\sigma})$ satisfy the product rule

$$\pi(R_{\lambda}^{\sigma}) \pi(R_{\eta}^{\sigma}) = \sigma(\lambda, \eta) \pi(R_{\lambda + \eta}^{\sigma}),$$

for $\sigma(\lambda, \eta) = \exp(2\pi i u_{\theta} \lambda \wedge \eta)$, which is the product rule of the twisted group algebra $C^*(\Lambda, \sigma)$. □

We obtain in this way an isospectral noncommutative geometry given by the finitely summable spectral triple

$$(7.18) \quad (C^{\infty}(X_{\epsilon})_u, L^2(X_{\epsilon}, S), \not{D}_{X_{\epsilon}}).$$

Corollary 7.4. *In the case $u = \theta$ and $u = \theta'$, the isospectral deformation (7.18) defines a finitely summable spectral triple for the noncommutative tori $\mathbb{T}_{\Lambda, i}$, with dense subalgebra $\mathbb{C}(\Lambda, \sigma)$.*

Proof. This is a direct consequence of Proposition 7.3 and the identifications of Corollary 4.2 of the $\mathbb{T}_{\Lambda, i}$ with twisted group C^* -algebras $C^*(\Lambda, \sigma)$. □

The representation (7.17) of $C^*(\Lambda, \sigma)$ extends to an action of $C^*(S(\Lambda, V), \tilde{\sigma})$, as follows. Let $U(k \log \epsilon)$ denote the unitary operator

$$(7.19) \quad (U(k \log \epsilon)\psi)((x, y), t) = \psi(A_{\epsilon}^k(x, y), t) = \psi((x, y), t - k \log \epsilon).$$

Proposition 7.5. *The operators*

$$(7.20) \quad \pi(R_{(\lambda,k)}^{\tilde{\sigma}}) := E_{\lambda} \Xi_u(\lambda, L_1, L_2) U(k \log \epsilon)$$

define a representation on $\mathcal{H} = L^2(X_{\epsilon}, S)$ of the twisted group C^* -algebra $C^*(\Lambda \rtimes_{\epsilon} V, \tilde{\sigma})$, for the cocycle

$$\tilde{\sigma}((\lambda, k), (\eta, r)) = \exp\left(2\pi i \frac{u_{\theta}}{2} \lambda \wedge A_{\epsilon}^k(\eta)\right).$$

Proof. We have the identities

$$\begin{aligned} (U(k \log \epsilon) E_{\eta}) &= E_{A_{\epsilon}^k(\eta)}, \\ \Xi_u(\lambda, L_1, L_2) E_{A_{\epsilon}^k(\eta)} &= e^{2\pi i u \lambda \wedge A_{\epsilon}^k(\eta)} E_{A_{\epsilon}^k(\eta)}, \\ E_{\lambda} E_{A_{\epsilon}^k(\eta)} &= E_{\lambda + A_{\epsilon}^k(\eta)}, \\ U(k \log \epsilon) \Xi_u(\eta, L_1, L_2) U(r \log \epsilon) &= \Xi_u(A_{\epsilon}^k(\eta), L_1, L_2) U((k+r) \log \epsilon). \end{aligned}$$

These combine to give the composition rule

$$\begin{aligned} E_{\lambda} \Xi_u(\lambda, L_1, L_2) U(k \log \epsilon) E_{\eta} \Xi_u(\eta, L_1, L_2) U(r \log \epsilon) = \\ \tilde{\sigma}((\eta, r), (\lambda, k)) E_{\lambda + A_{\epsilon}^k(\eta)} \Xi_u(\lambda + A_{\epsilon}^k(\eta), L_1, L_2) U((k+r) \log \epsilon). \end{aligned}$$

□

7.3. Unitary equivalences. We begin by reformulating the data described above in an equivalent form by expanding in Fourier modes along the fiber tori as in [2]. Recall that the fiber over $t \in [0, \log \epsilon)$ is given by the torus $T_t^2 = \mathbb{R}^2 / \Lambda_t$, with $\Lambda_t = \Theta_t(\Lambda)$. Thus, if we denote by (x, y) , as above, the coordinates in T_t^2 , we can write these as $(x, y) = \Theta_t(a, b)$, with $(a, b) \in T^2 = \mathbb{R}^2 / \Lambda$, the reference torus. This means writing the spinors $\psi((x, y), t)$ in the form

$$(7.21) \quad \sum_{\lambda} \psi_{\lambda} e^{2\pi i \langle (a,b), \lambda \rangle} = \sum_{\lambda} \psi_{\lambda} e^{2\pi i \langle \Theta_{-t}(x,y), \lambda \rangle} = \sum_{\lambda} \psi_{\lambda} E_{\lambda}.$$

The Dirac operator acts on E_{λ} as

$$\hat{\partial}_{X_{\epsilon}} E_{\lambda} = \left(\frac{\partial}{\partial t} \sigma_0 + 2\pi i \lambda_1 \sigma_1 + 2\pi i \lambda_2 \sigma_2 \right) E_{\lambda}.$$

The operators $\pi(R_{\eta}^{\sigma})$ act as

$$(7.22) \quad E_{\eta} \Xi_u(\eta, L_1, L_2) E_{\lambda} = e^{2\pi i u \eta \wedge \lambda} E_{\eta + \lambda}.$$

The commutators are bounded operators of the form

$$(7.23) \quad [\hat{\partial}_{X_{\epsilon}}, \pi(R_{\eta}^{\sigma})] = (\eta_1 \sigma_1 + \eta_2 \sigma_2) R_{\eta}^{\sigma}.$$

Thus, passing to Fourier modes in the fiber directions gives a unitarily equivalent spectral triple for the noncommutative tori $\mathbb{T}_{\Lambda, i}$, with

$$(7.24) \quad \begin{aligned} \hat{\partial}_{X_{\epsilon}} \psi_{\lambda} &= \left(\frac{\partial}{\partial t} \sigma_0 + 2\pi i \lambda_1 \sigma_1 + 2\pi i \lambda_2 \sigma_2 \right) \psi_{\lambda} \\ \hat{\pi}(R_{\eta}^{\sigma}) \psi_{\lambda} &= \sigma(\eta, \lambda) \psi_{\lambda + \eta}. \end{aligned}$$

We then consider a second unitary equivalence, which, as in [2] adjusts for the possible signs of λ_1 and λ_2 . Namely, we define the following unitary operator on the Hilbert space of the spinors ψ_λ . We set

$$(7.25) \quad \mathcal{U} \psi_\lambda = \sigma_\lambda \psi_\lambda,$$

where σ_λ is a product of Pauli matrices, where σ_i , for $i = 1, 2$ appears in the product if and only if $\lambda_i < 0$. Then the Dirac operator transforms to the unitarily equivalent operator

$$(7.26) \quad \mathcal{U} \hat{\mathcal{D}}_{X_\epsilon} \mathcal{U}^* = \text{sign}(N(\lambda)) \left(\frac{\partial}{\partial t} \sigma_0 + 2\pi i |\lambda_1| \sigma_1 + 2\pi i |\lambda_2| \sigma_2 \right).$$

The action of the R_η^σ transform correspondingly to the operators

$$(7.27) \quad \mathcal{U} \hat{\pi}(R_\eta^\sigma) \mathcal{U}^* : \sigma_\lambda \psi_\lambda \mapsto \sigma_{\lambda+\eta} \psi_{\lambda+\eta}.$$

We then perform the other unitary transformation used in [2]. To this purpose, let us fix a choice of a fundamental domain \mathcal{F}_V for the action of V on the lattice Λ . By this choice of a fundamental domain, we can write uniquely an element $\lambda \in \Lambda$ in the form $\lambda = A_\epsilon^k(\mu)$, for a $\mu \in \mathcal{F}_V$ and a $k \in \mathbb{Z}$.

For $\lambda = A_\epsilon^k(\mu) \neq 0$, consider then the time shift

$$(7.28) \quad \tilde{\mathcal{U}}(\sigma_\lambda \psi_\lambda)(t) = \sigma_\lambda \psi_\lambda(t - \log \frac{|\mu_1|}{|N(\mu)|^{1/2}}),$$

so that we have

$$(7.29) \quad \tilde{\psi}_\lambda := \tilde{\mathcal{U}}(\sigma_\lambda \psi_\lambda) = \sigma_\lambda \psi_{|N(\lambda)|^{1/2}(\text{sign}(\lambda_1)\epsilon^k, \text{sign}(\lambda_2)\epsilon^{-k})}.$$

One obtains in this way a unitarily equivalent spectral triple for $\mathbb{T}_{\Lambda,i}$, with Dirac operator

$$(7.30) \quad \tilde{\mathcal{D}} = \tilde{\mathcal{D}}^{(0)} + \sum_{\mu \in (\Lambda \setminus \{0\})/V} \tilde{\mathcal{D}}^{(\mu)},$$

where

$$(7.31) \quad \begin{aligned} \tilde{\mathcal{D}}^{(\mu)} \tilde{\psi}_{A_\epsilon^k(\mu)} = \\ \text{sign}(N(\mu)) |N(\mu)|^{1/2} \left(|N(\mu)|^{-1/2} \frac{\partial}{\partial t} \sigma_0 + 2\pi i \epsilon^k \sigma_1 + 2\pi i \epsilon^{-k} \sigma_2 \right) \tilde{\psi}_{A_\epsilon^k(\mu)} \end{aligned}$$

while the action of the R_η^σ is by

$$\tilde{\pi}(R_\eta^\sigma) \tilde{\psi}_\lambda = \tilde{\psi}_{\lambda+\eta}.$$

As in [2], one can write the operator $\tilde{\mathcal{D}}^{(\mu)}$ as a product

$$\tilde{\mathcal{D}}^{(\mu)} = D_\mu B_\mu,$$

with

$$(7.32) \quad \begin{aligned} D_\mu \tilde{\psi}_{A_\epsilon^k(\mu)} &= \text{sign}(N(\mu)) |N(\mu)|^{1/2} \tilde{\psi}_{A_\epsilon^k(\mu)} \\ B_\mu \tilde{\psi}_{A_\epsilon^k(\mu)} &= (|N(\mu)|^{-1/2} \frac{\partial}{\partial t} \sigma_0 + 2\pi i \epsilon^k \sigma_1 + 2\pi i \epsilon^{-k} \sigma_2) \tilde{\psi}_{A_\epsilon^k(\mu)}. \end{aligned}$$

In the following, we relate the Dirac operator $\hat{\mathcal{D}}_{X_\epsilon}$, its unitarily equivalent operators discussed here above, and the decomposition (7.32) to known differential operators on noncommutative tori.

7.4. Differential operators on noncommutative tori. Notice that the action on \mathbb{R}^2 of the 1-paramater subgroup of $\mathrm{SL}_2(\mathbb{R})$

$$\Theta_t = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}$$

has fixed point $(0, 0)$, with stable manifold the axis $(0, y)$ and unstable manifold the axis $(x, 0)$. On the standard torus $\mathbb{R}^2/\mathbb{Z}^2$ with coordinates (s_1, s_2) with $(x, y) = (s_1 + s_2\theta, s_1 + s_2\theta')$ these two directions define the two Kronecker foliations $s_1 + s_2\theta$ and $s_1 + s_2\theta'$ with conjugate slopes θ and θ' . The points of the lattice Λ determine on these two foliations the points of the pseudolattices $\mathbb{Z} + \mathbb{Z}\theta$ and $\mathbb{Z} + \mathbb{Z}\theta'$, which define the equivalence relation on the space of leaves of the two Kronecker foliations, defining as quotients the noncommutative tori $\mathbb{T}_{\Lambda, i}$, $i = 1, 2$. The action of Θ_t is expanding along the line $L_\theta = \{s_1 + s_2\theta\}$ and contracting along $L_{\theta'} = \{s_1 + s_2\theta'\}$ and flows the other points of \mathbb{R}^2 along hyperbola with asymptotes $L_{\theta'}$ and L_θ .

Thus, the operators $e^t \frac{\partial}{\partial x}$ and $e^{-t} \frac{\partial}{\partial y}$ correspond to derivations along the leaf direction of these two transverse Kronecker foliations. The factors e^t and e^{-t} are the normalization factors that account for the rescaling of the transverse measure due to the action of the flow Θ_t . In fact, consider for instance a small transversal of length ℓ for the Kronecker foliation L_θ , given by the interval $T_\ell = \{(x, y) : x = 1, -\ell/2 < y < \ell/2\}$. The flow Θ_t maps it to the transversal $\Theta_t(T_\ell) = \{(x = e^t, y) : -e^{-t}\ell/2 < y < e^{-t}\ell/2\}$ of length $e^{-t}\ell$. Thus, the differentiation $\frac{\partial}{\partial x}$ in the leaf direction of L_θ is weighted by the factor e^t that normalizes the length of the transversal and corrects for the scaling of the transverse measure.

Consider then the terms $2\pi i\lambda_1\sigma_1$ and $2\pi i\lambda_2\sigma_2$ in the operator

$$\hat{\mathcal{D}}_{X_\epsilon} : \psi_\lambda \mapsto \left(\frac{\partial}{\partial t} \sigma_0 + 2\pi i\lambda_1\sigma_1 + 2\pi i\lambda_2\sigma_2 \right) \psi_\lambda,$$

that we obtained after passing to Fourier modes on the fiber tori T_t^2 . These terms correspond, respectively, to the leafwise derivations $e^t \frac{\partial}{\partial x}$ and $e^{-t} \frac{\partial}{\partial y}$. These can be expressed equivalently in terms of the operators

$$\delta_\theta : \psi_{n,m} \mapsto (n + m\theta) \psi_{n,m}, \quad \text{and} \quad \delta_{\theta'} : \psi_{n,m} \mapsto (n + m\theta') \psi_{n,m},$$

so that the sum $\lambda_1\sigma_1 + \lambda_2\sigma_2$ acts as the operator

$$(7.33) \quad \mathcal{D}_{\theta, \theta'} = \begin{pmatrix} 0 & \delta_{\theta'} - i\delta_\theta \\ \delta_{\theta'} + i\delta_\theta & 0 \end{pmatrix}.$$

This gives the Dirac operator of a spectral triple on the noncommutative tori $\mathbb{T}_{\Lambda, i}$ with

$$R_{r,k}^\sigma \psi_{n,m} = \sigma((r, k), (n, m)) \psi_{(n,m)+(r,k)}$$

and

$$[\mathcal{D}_{\theta, \theta'}, R_{r,k}^\sigma] = \begin{pmatrix} 0 & (r + k\theta') - i(r + k\theta) \\ (r + k\theta') + i(r + k\theta) & 0 \end{pmatrix} R_{r,k}^\sigma.$$

In the particular case where $\theta' = -\theta$, this agrees with the spectral triple for the first order signature operator on the noncommutative torus considered, for instance, in [21]. The construction of [21] can be interpreted as obtained by using the two transverse Kronecker foliations L_θ and $L_{-\theta}$ and the associated leafwise derivations $\partial/\partial x$ and $\partial/\partial y$.

We can consider here the same kind of unitary transformations that we described earlier for $\hat{\mathcal{D}}_{X_\epsilon}$, applied to the operator $\mathcal{D}_{\theta,\theta'}$ of (7.33). Let us denote by $\mathcal{D}_{\theta,\theta',0}$ the restriction of $\mathcal{D}_{\theta,\theta'}$ to the complement of the zero modes ψ_0 (*i.e.* $\lambda = 0$). We have, as in (7.30),

$$(7.34) \quad \mathcal{D}_{\theta,\theta',0} = \sum_{\mu \in (\Lambda \setminus \{0\})/V} \mathcal{D}_{\theta,\theta'}^\mu,$$

with

$$\mathcal{D}_{\theta,\theta'}^\mu \psi_{A_\epsilon^k(\mu)} = (\lambda_1 \sigma_1 + \lambda_2 \sigma_2) \psi_{A_\epsilon^k(\mu)}.$$

After the unitary transformation $\tilde{\mathcal{U}}\mathcal{U}$ with \mathcal{U} as in (7.25) and $\tilde{\mathcal{U}}$ as in (7.29), we obtain a unitarily equivalent operator

$$(7.35) \quad \tilde{\mathcal{D}}_{\theta,\theta'}^\mu \tilde{\psi}_{A_\epsilon^k(\mu)} = \text{sign}(N(\mu)) |N(\mu)|^{1/2} (\epsilon^k \sigma_1 + \epsilon^{-k} \sigma_2) \tilde{\psi}_{A_\epsilon^k(\mu)}.$$

As before, we factor this as a product of the operators

$$(7.36) \quad \tilde{\mathcal{D}}_{\theta,\theta'}^\mu = D_\theta^\mu B_\theta,$$

with

$$(7.37) \quad \begin{aligned} D_\theta^\mu \tilde{\psi}_{A_\epsilon^k(\mu)} &= \text{sign}(N(\mu)) |N(\mu)|^{1/2} \tilde{\psi}_{A_\epsilon^k(\mu)} \\ B_\theta \tilde{\psi}_{A_\epsilon^k(\mu)} &= (\epsilon^k \sigma_1 + \epsilon^{-k} \sigma_2) \tilde{\psi}_{A_\epsilon^k(\mu)}. \end{aligned}$$

8. SHIMIZU L -FUNCTION AND LORENTZIAN GEOMETRY

In this section we describe another way of relating the Shimizu L -function to the geometry of noncommutative tori with real multiplication, by regarding the norms $N(\lambda)$, for $\lambda \in \Lambda$, as defining the momenta of a *Lorentzian* rather than Euclidean Dirac operator.

Instead of working with positive inner product spaces, as in the case of Euclidean spectral triples, the Galois involution of the real quadratic field defines a natural choice of a “Krein involution” and the norm correspondingly defines an indefinite quadratic form. One formulates in this way a notion of spectral triple over a real quadratic field and with Lorentzian signature, using the relation between indefinite inner product spaces and the associated real Hilbert spaces. The main point that requires care is the fact that the Lorentzian Dirac operator has a noncompact group of symmetries, in our case given by the units of the real quadratic field, hence it fails to have compact resolvent due to the presence of infinite multiplicities in the eigenvalues. We show that the multiplicities can be resolved by transforming the triple via a Krein isometry that is an unbounded self-adjoint operator in the associated real Hilbert space and defines a finitely summable associated Dirac operator in the Euclidean signature.

As above, we let Λ be the lattice in \mathbb{R}^2 associated to a lattice $L \subset \mathbb{K}$ in a real quadratic field $\mathbb{K} = \mathbb{Q}(\sqrt{d})$ by the embeddings $\iota_i : \mathbb{K} \hookrightarrow \mathbb{R}$,

$$(8.1) \quad \Lambda = \{\lambda \in \mathbb{R}^2 \mid \lambda = (\lambda_1, \lambda_2) = (\iota_1(\ell), \iota_2(\ell)), \ell \in L\}.$$

We denote, as above, by V the group $V = \epsilon^{\mathbb{Z}}$ of units preserving Λ . We denote the action as above with $\lambda \mapsto A_\epsilon^k(\lambda) = (\epsilon^k \lambda_1, \epsilon^{-k} \lambda_2)$.

For $x \in \mathbb{K}$, we denote by $x' = c(x)$ the image under the Galois involution of \mathbb{K} . For $\lambda = (\lambda_1, \lambda_2) \in \Lambda$, we have $\lambda_2 = c(\lambda_1)$. The norm is given by $N(\lambda) = \lambda_1 \lambda_2$, and $N(\epsilon) = \epsilon \epsilon' = 1$.

We consider the quadratic form $N(\lambda) = \lambda_1 \lambda_2 = (n + m\theta)(n + m\theta')$ to be the analog of the wave operator $\square = p_0^2 - p_1^2$.

Its Dirac factorization into linear first order operator is obtained by considering a linear operator of the form

$$(8.2) \quad \mathcal{D}_\lambda = \begin{pmatrix} 0 & \mathcal{D}_\lambda^+ \\ \mathcal{D}_\lambda^- & 0 \end{pmatrix} := \begin{pmatrix} 0 & \lambda_1 \\ \lambda_2 & 0 \end{pmatrix},$$

whose square is $\mathcal{D}_\lambda^2 = \square_\lambda$, with

$$(8.3) \quad \square_\lambda = \begin{pmatrix} N(\lambda) & 0 \\ 0 & N(\lambda) \end{pmatrix}.$$

We assemble these modes to define an operator \mathcal{D} acting on $\mathcal{H} = \ell^2(\Lambda) \oplus \ell^2(\Lambda)$ by $\mathcal{D}e_{\lambda,\pm} = \mathcal{D}_\lambda e_{\lambda,\pm}$. This satisfies $\mathcal{D}\gamma = -\gamma\mathcal{D}$ with respect to the $\mathbb{Z}/2\mathbb{Z}$ -grading

$$\gamma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Consider the algebra $C^*(\Lambda, \sigma)$ of the noncommutative torus acting diagonally on \mathcal{H} . The operator \mathcal{D} has bounded commutators with the elements of the dense subalgebra $\mathbb{C}(\Lambda, \sigma)$ since we have

$$(8.4) \quad [\mathcal{D}, R_\eta^\sigma]e_{\lambda,\pm} = \sigma(\lambda, \eta) \eta_\mp e_{\eta+\lambda,\pm},$$

where we used the notation $\eta_+ = \eta_1$ and $\eta_- = \eta_2$.

However, the other properties of \mathcal{D} differ significantly from what one usually postulates for Dirac operators of spectral triples.

First of all, notice that \mathcal{D} is not self-adjoint. In fact, it is invariant with respect to a different involution, defined for operators with coefficients in the real quadratic field \mathbb{K} , namely

$$\mathcal{D} = c(\mathcal{D}^t),$$

where $\mathcal{D}^t = (\mathcal{D}_\lambda^t)$ denotes the transpose and $c(\mathcal{D})$ denotes the effect of the Galois involution $c : x \mapsto x'$ of \mathbb{K} applied to the coefficients of \mathcal{D} . In this arithmetic context, it is natural to require this property instead of self-adjointness.

A more serious problem, however, comes from the fact that the operator \mathcal{D} has infinite multiplicities, hence it is very far from having the compact resolvent property of spectral triple. This is a typical problem one encounters in trying to extend the formalism of spectral triples from the Euclidean to the Lorentzian context, because of the presence of non-compact symmetry groups for Lorentzian manifold. Here the non-compact symmetry group is given by the units in $V = \epsilon^{\mathbb{Z}}$.

8.1. Arithmetic Krein spaces. It is well known that, when one replaces Euclidean geometry by Lorentzian geometry, the notion of Hilbert space is replaced by the notion of a Krein space (cf. e.g. [7]). The version we consider here is slightly different from the usual one, since we want to be able to work over the real quadratic field \mathbb{K} instead of passing directly to complex numbers.

Definition 8.1. Let $c : \mathbb{K} \rightarrow \mathbb{K}$ denote the Galois involution $c : x \mapsto x'$ of the real quadratic field. Let \mathcal{V} be a \mathbb{K} -vector space. We say that a map $T : \mathcal{V} \rightarrow \mathcal{V}$ is c -linear if it satisfies $T(av + bw) = c(a)T(v) + c(b)T(w)$.

A Lorentzian pairing on a \mathbb{K} -vector space \mathcal{V} is a non-degenerate \mathbb{K} -valued pairing

$$(\cdot, \cdot) : \mathcal{V} \times \mathcal{V} \rightarrow \mathbb{K}$$

which is c -linear in the first variable and linear in the second.

We can then introduce the analog of the notion of a Krein space, in this arithmetic context.

Definition 8.2. A Krein space over a real quadratic field \mathbb{K} (or \mathbb{K} -Krein space) is a \mathbb{K} -vector space \mathcal{V} endowed with a Lorentzian pairing (\cdot, \cdot) as in Definition 8.1, and a c -linear involution $\kappa : \mathcal{V} \rightarrow \mathcal{V}$, such that the pairing $(\kappa \cdot, \cdot)$ has the following properties:

- (1) $(\kappa \cdot, \cdot) = c(\cdot, \kappa \cdot)$
- (2) For all $v \neq 0$ in \mathcal{V} , the elements $(\kappa v, v) \in \mathbb{K}$ are totally positive.

When properties (1) and (2) of Definition 8.2 holds, we say that $(\kappa \cdot, \cdot)$ defines a positive definite inner product. We have a corresponding notion of Krein adjoint as follows.

Definition 8.3. Given a \mathbb{K} -linear operator T on a \mathbb{K} -Krein space \mathcal{V} , the Krein adjoint T^\dagger is the adjoint in the Lorentzian pairing (\cdot, \cdot) ,

$$(8.5) \quad (v, Tw) = (T^\dagger v, w).$$

The c -linear involution κ of Definition 8.2 corresponds to a Wick rotation from Lorentzian to Euclidean signature. The Krein adjoint satisfies $T^\dagger = \kappa T^* \kappa$, where T^* is the adjoint in the inner product $\langle \cdot, \cdot \rangle = (\kappa \cdot, \cdot)$.

Given a \mathbb{K} -Krein space \mathcal{V} , there are two naturally associated real Hilbert spaces, obtained by considering the real vector spaces

$$(8.6) \quad \mathcal{V}_{\mathbb{R}, i} := \mathcal{V} \otimes_{\iota_i(\mathbb{K})} \mathbb{R},$$

obtained by tensoring \mathcal{V} with \mathbb{R} using either one of the two embeddings $\iota_i : \mathbb{K} \hookrightarrow \mathbb{R}$ of the real quadratic field.

Lemma 8.4. The pairing

$$(8.7) \quad \langle v, w \rangle = \frac{1}{2} \iota_1((\kappa v, w) + (v, \kappa w)) = \frac{1}{2} \iota_2((\kappa v, w) + (v, \kappa w))$$

induced on $\mathcal{V}_{\mathbb{R}, i}$ by the Lorentzian pairing (\cdot, \cdot) on \mathcal{V} defines a real valued positive definite inner product.

Proof. We know that $(v, \kappa w) = c(\kappa v, w)$. Thus, we have

$$\langle v, w \rangle = \frac{1}{2} (\iota_1(\kappa v, w) + \iota_2(\kappa v, w)).$$

We can extend this pairing by \mathbb{R} -linearity to define a bilinear form on $\mathcal{V}_{\mathbb{R},i}$. By the assumption that for $v \neq 0$ the $(\kappa v, v)$ are totally positive elements of \mathbb{K} we obtain that (8.7) defines a positive-definite inner product. \square

in the following, we still denote by $\mathcal{V}_{\mathbb{R},i}$ the Hilbert space completion obtained in this way.

8.2. Lorentzian spectral triples over real quadratic fields. It is not uncommon to make use of Krein spaces to extend the formalism of spectral triples to Lorentzian geometry, see *e.g.* [28]. Here we follow a similar viewpoint, adapted to the arithmetic setting of real quadratic fields.

For a \mathbb{K} -linear operator T acting on a \mathbb{K} -Krein space \mathcal{V} , we define $\mathbb{M}_i(T) \geq -\infty$, for $i = 1, 2$, as

$$(8.8) \quad \mathbb{M}_i(T) := \inf_{(v,v)=1} \iota_i(Tv, Tv).$$

We introduce the following preliminary notion of a \mathbb{K} -triple, which we then refine by additional properties providing the analog of a spectral triple.

Definition 8.5. *A Krein \mathbb{K} -triple consists of data $(\mathcal{A}, \mathcal{V}, \mathcal{D})$ with the following properties.*

- (1) \mathcal{A} is an involutive algebra over the real quadratic field \mathbb{K} .
- (2) \mathcal{V} is a \mathbb{K} -Krein space with non-degenerate \mathbb{K} -bilinear form (\cdot, \cdot) .
- (3) The algebra \mathcal{A} acts on \mathcal{V} via a representation $\pi : \mathcal{A} \rightarrow \text{End}_{\mathbb{K}}(\mathcal{V})$, with the involution of \mathcal{A} realized by the Krein adjoint $\pi(a^*) = \pi(a)^\dagger$.
- (4) The operators $\pi(a)$, for $a \in \mathcal{A}$, satisfy

$$(8.9) \quad \mathbb{M}_i(a) > -\infty.$$

- (5) The operator \mathcal{D} is a densely defined \mathbb{K} -linear operator on \mathcal{V} , which is Krein-self-adjoint, $\mathcal{D}^\dagger = \mathcal{D}$.
- (6) The commutators $C_a := [\mathcal{D}, a]$ satisfy

$$(8.10) \quad \mathbb{M}_i(C_a) > -\infty, \quad \forall a \in \mathcal{A}.$$

We then define Lorentzian \mathbb{K} -spectral triples in the following way.

Definition 8.6. *A Krein \mathbb{K} -triple $(\mathcal{A}, \mathcal{V}, \mathcal{D})$ as in Definition 8.5 is a finitely summable Lorentzian \mathbb{K} -spectral triple if the following holds.*

- (1) There exists a densely defined \mathbb{K} -linear operator $U : \mathcal{V} \rightarrow \mathcal{V}$ with $(Uv, Uv) = (v, v)$, for all $v \in \text{Dom}(U)$ and $U^\dagger = U^{-1}$, with the property that

$$(8.11) \quad U^\dagger \mathcal{D} U = \mathcal{D}.$$

- (2) The commutators $C_{a,U} := [\mathcal{D}_U, \pi_U(a)]$, with $\pi_U(a) = U^\dagger \pi(a) U$, satisfy the condition

$$(8.12) \quad \mathbb{M}_i(C_{a,U}) > -\infty, \quad \forall a \in \mathcal{A}.$$

(3) The operator U is an unbounded self-adjoint operator, $U = U^*$ on the associated real Hilbert space $\mathcal{V}_{\mathbb{R},i}$ with the inner product $\langle \cdot, \cdot \rangle$ of (8.7).

(4) The triple $(\mathcal{A}, \mathcal{V}, \mathcal{D})$ is p -summable for $p \in \mathbb{R}_+^*$ if

$$(8.13) \quad \sum_n |\langle Ue_n, |\mathcal{D}^2|Ue_n \rangle|^{-s/2} < \infty, \quad \forall s \geq p,$$

where e_n is an orthonormal basis for the complement of the zero modes of the operator $|\mathcal{D}^2|$ in the real Hilbert space $\mathcal{V}_{\mathbb{R},i}$.

Notice that in Krein spaces isometries are not necessarily bounded operators (see [7] §VI), so the U is only densely defined in general. The definition given here is different from the notions of Lorentzian spectral triples currently developed in the literature. The differences stem mainly from our need to work over a finite extension of \mathbb{Q} instead of \mathbb{C} and to resolve the infinite multiplicity of the eigenvalues. We also require the weaker property (8.9), (8.10) and (8.12), instead of requiring continuity in the operator norm in the associated Hilbert space. These conditions will become more transparent in our main example below.

8.3. Arithmetic twisted group algebras. We consider the \mathbb{K} -vector space \mathcal{V}_Λ spanned by the basis elements e_λ with $\lambda \in \Lambda$, endowed with the pairing

$$(8.14) \quad (v, w) := \sum_\lambda c(a_\lambda) b_\lambda,$$

for $v = \sum_\lambda a_\lambda e_\lambda$ and $w = \sum_\lambda b_\lambda e_\lambda$, and with $c : x \mapsto x'$ the Galois involution of \mathbb{K} .

Lemma 8.7. *The space \mathcal{V}_Λ with the pairing (8.14) is a \mathbb{K} -Krein space.*

Proof. Clearly the pairing (8.14) is a Lorentzian pairing in the sense of Definition 8.1. Let $\kappa : \mathcal{V}_\Lambda \rightarrow \mathcal{V}_\Lambda$ be given by the Galois involution

$$\kappa(v) = \sum_\lambda c(a_\lambda) e_\lambda.$$

Then the pairing $\langle v, w \rangle = (\kappa v, w) = c(v, \kappa w)$ is a positive definite inner product, as in Definition 8.2. In fact, we have

$$\iota_1 \langle v, v \rangle = \sum_\lambda \iota_1(a_\lambda^2) \geq 0, \quad \iota_2 \langle v, v \rangle = \sum_\lambda \iota_2(a_\lambda^2) \geq 0.$$

□

We consider on \mathcal{V}_Λ the action of the group ring $\mathbb{K}[\Lambda]$, given by $R_\lambda e_\eta = e_{\lambda+\eta}$.

Lemma 8.8. *The operators R_λ acting on \mathcal{V}_Λ satisfy*

$$\mathbb{M}_i(R_\lambda) > -\infty.$$

Moreover, the operators R_λ define bounded operators in the associated real Hilbert spaces $\mathcal{V}_{\Lambda, \mathbb{R}, i} = \mathcal{V}_\Lambda \otimes_{\iota_i(\mathbb{K})} \mathbb{R}$.

Proof. The operators R_λ are Krein isometries, and $(R_\lambda v, R_\lambda v) = (v, v)$ implies that $\mathbb{M}_i(R_\lambda) = 1$. The operators R_λ act by $e_\eta \mapsto e_{\lambda+\eta}$ on the associated Hilbert spaces, hence they define bounded (unitary) operators. □

Now we want to introduce, in this setting of \mathbb{K} -Krein spaces, an analog of the twisted group ring $\mathbb{C}(\Lambda, \sigma)$ (the noncommutative torus) we have been working with in the complex case.

Lemma 8.9. *Suppose given $\omega \in \mathbb{K}^*$ with $N(\omega) = \omega\omega' = 1$. Then the expression*

$$(8.15) \quad \varpi(\lambda, \eta) = \omega^{(n,m) \wedge (r,k)},$$

for $\lambda = (n + m\theta, n + m\theta')$ and $\eta = (r + k\theta, r + k\theta')$, defines a \mathbb{K}^* -valued group 2-cocycle σ on Λ .

Proof. The argument is the same as in the complex case. It suffices to show that the cocycle condition holds. \square

Definition 8.10. *The twisted group ring $\mathbb{K}(\Lambda, \sigma)$ is the unital involutive \mathbb{K} -algebra generated by elements R_λ^ϖ with the product*

$$(8.16) \quad R_\lambda^\varpi R_\eta^\varpi = \varpi(\lambda, \eta) R_{\lambda+\eta}^\varpi = \omega^{(n,m) \wedge (r,k)} R_{\lambda+\eta}^\varpi,$$

for $\lambda = (n + m\theta, n + m\theta')$ and $\eta = (r + k\theta, r + k\theta')$, and the involution $(R_\lambda^\varpi)^* = R_{-\lambda}^\varpi$.

The twisted group ring $\mathbb{K}(\Lambda, \sigma)$ also acts on \mathcal{V}_Λ by

$$(8.17) \quad R_\lambda^\varpi e_\eta = \varpi(\eta, \lambda) e_{\lambda+\eta}.$$

Lemma 8.11. *The operators R_λ^ϖ , acting as in (8.17), satisfy $\mathbb{M}_i(R_\lambda^\varpi) > -\infty$.*

Proof. The action (8.17) preserves the Lorentzian pairing (\cdot, \cdot) on \mathcal{V}_Λ since

$$(R_\lambda^\varpi e_\eta, R_\lambda^\varpi e_\zeta) = c(\omega(\eta, \lambda)) \omega(\zeta, \lambda) \delta_{\eta, \zeta} = N(\omega(\eta, \lambda)) \delta_{\eta, \zeta} = (e_\eta, e_\zeta),$$

since $N(\omega(\eta, \lambda)) = 1$. The condition $(R^\varpi v, R^\varpi v) = (v, v)$ implies $\mathbb{M}_i(R_\lambda^\varpi) = 1$. \square

However, notice that, while the action of $\mathbb{K}[\Lambda]$ extends to an action by bounded operators on the associated real Hilbert spaces $\mathcal{V}_{\mathbb{K}} \otimes_{\iota_i(\mathbb{K})} \mathbb{R}$, the induced action on $\mathcal{V}_{\mathbb{R}}$ of the twisted group ring $\mathbb{K}(\Lambda, \varpi)$ is by the unbounded operators

$$(8.18) \quad R_\lambda^\varpi e_{\eta, \pm} = A_\omega^{(r,k) \wedge (n,m)} e_{\lambda+\eta, \pm},$$

with

$$(8.19) \quad A_\omega = \begin{pmatrix} \iota_1(\omega) & 0 \\ 0 & \iota_2(\omega) \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R}).$$

As in the complex case, we can also consider the group ring $\mathbb{K}[S(\Lambda, V)]$ for $S(\Lambda, V) = \Lambda \rtimes V$. The cocycle (8.15) extends to a cocycle on the cross product by setting

$$(8.20) \quad \tilde{\varpi}((\lambda, k), (\eta, r)) = \varpi(\lambda, A_\epsilon^k(\eta)) = \omega^{(n,m) \wedge (u,v) \varphi_\epsilon^k},$$

for $\lambda = (n + m\theta, n + m\theta')$ and $\eta = (u + v\theta, u + v\theta')$, with $n, m, u, v \in \mathbb{Z}$.

Definition 8.12. *The twisted group ring $\mathbb{K}(S(\Lambda, V), \tilde{\varpi})$ is the unitary involutive \mathbb{K} -algebra with generators $R_{\lambda, k}^{\tilde{\varpi}}$ satisfying*

$$R_{\lambda, k}^{\tilde{\varpi}} R_{\eta, r}^{\tilde{\varpi}} = \tilde{\varpi}((\lambda, k), (\eta, r)) R_{\lambda + A_\epsilon^k(\eta), k+r}^{\tilde{\varpi}},$$

with the involution $(R_{\lambda, k}^{\tilde{\varpi}})^* = R_{-A_\epsilon^{-k}(\lambda), -k}^{\tilde{\varpi}}$.

8.4. Lorentzian Dirac operator. On the \mathbb{K} -Krein space $\mathcal{V}_\Lambda \oplus \mathcal{V}_\Lambda$ we consider the densely defined \mathbb{K} -linear operator

$$(8.21) \quad \mathcal{D}_\mathbb{K} e_{\lambda, \pm} = \mathcal{D}_{\mathbb{K}, \lambda} e_{\lambda, \pm} = \begin{pmatrix} 0 & \mathcal{D}_\lambda^+ \\ \mathcal{D}_\lambda^- & 0 \end{pmatrix} e_{\lambda, \pm} := \begin{pmatrix} 0 & \ell \\ c(\ell) & 0 \end{pmatrix} e_{\lambda, \pm},$$

where we write $\lambda \in \Lambda$ as $\lambda = (\iota_1(\ell), \iota_2(\ell))$ with $\ell \in L \subset \mathbb{K}$, as in (8.1).

The operator $\mathcal{D}_\mathbb{K}$ of (8.21) induces on the real Hilbert space $\mathcal{V}_{\mathbb{R}, i} \oplus \mathcal{V}_{\mathbb{R}, i}$ the \mathbb{R} -linear operators

$$(8.22) \quad \mathcal{D}_\lambda = \begin{pmatrix} 0 & \lambda_1 \\ \lambda_2 & 0 \end{pmatrix} \quad \text{and} \quad c(\mathcal{D}_\lambda) = \begin{pmatrix} 0 & \lambda_2 \\ \lambda_1 & 0 \end{pmatrix},$$

respectively. This recovers the Lorentzian Dirac operator described in (8.2) above.

Lemma 8.13. *The data $(\mathbb{K}(\Lambda, \varpi), \mathcal{V}_\Lambda \oplus \mathcal{V}_\Lambda, \mathcal{D}_\mathbb{K})$ define a Krein \mathbb{K} -triple in the sense of Definition 8.5.*

Proof. Properties (1)–(4) of Definition 8.5 follow from Lemma 8.7, Lemma 8.11, and the fact that the Krein adjoint $(R_\lambda^\varpi)^\dagger = R_{-\lambda}^\varpi = (R_\lambda^\varpi)^{-1}$. Property (5) follows directly from (8.21), since

$$\mathcal{D}_\mathbb{K}^\dagger = c(\mathcal{D}_{\mathbb{K}, \lambda}^t) = \begin{pmatrix} 0 & c(\lambda_2) \\ c(\lambda_1) & 0 \end{pmatrix} = \begin{pmatrix} 0 & \lambda_1 \\ \lambda_2 & 0 \end{pmatrix} = \mathcal{D}_\mathbb{K}.$$

We then need to prove (6), namely that the commutators $[\mathcal{D}_\mathbb{K}, R_\lambda^\varpi]$ satisfy

$$\mathbb{M}_i([\mathcal{D}_\mathbb{K}, R_\lambda^\varpi]) > -\infty.$$

We have

$$[\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] e_{\eta, \pm} = \varpi(\eta, \lambda) \begin{pmatrix} 0 & (\lambda_1 + \eta_1) - \eta_1 \\ (\lambda_2 + \eta_2) - \eta_2 & 0 \end{pmatrix} e_{\lambda + \eta, \pm}.$$

Thus, we have

$$([\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] v, [\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] v) = N(\lambda)(v, v),$$

from which the result follows. \square

Suppose given a choice of a fundamental domain \mathcal{F}_V for the action of $V = \epsilon^\mathbb{Z}$ on Λ .

Let $\rho(\lambda) \in \mathbb{Z}$ denote the unique integer such that $\lambda = A_\epsilon^{\rho(\lambda)}(\mu)$, with $\mu \in \mathcal{F}_V$.

Consider the \mathbb{K} -linear operator on $\mathcal{V}_\Lambda \oplus \mathcal{V}_\Lambda$ defined by

$$(8.23) \quad T_\epsilon e_{\lambda, \pm} := \begin{pmatrix} \epsilon^{\rho(\lambda)} & 0 \\ 0 & \epsilon^{-\rho(\lambda)} \end{pmatrix} e_{\lambda, \pm}.$$

Consider also the involution $J : \mathcal{V} \rightarrow \mathcal{V}$ defined by setting

$$J e_{\lambda, \pm} = e_{J(\lambda), \pm},$$

where $J(\lambda) = A_\epsilon^{-k}(\mu)$ for $\lambda = A_\epsilon^k(\mu)$ with $\mu \in \mathcal{F}_V$ and $k \in \mathbb{Z}$. This satisfies $J^2 = 1$ and $J^\dagger = J$.

We set $U_\epsilon = T_\epsilon J$, with

$$(8.24) \quad U_\epsilon e_{\lambda, \pm} = \begin{pmatrix} \epsilon^{-\rho(\lambda)} & 0 \\ 0 & \epsilon^{\rho(\lambda)} \end{pmatrix} e_{J(\lambda), \pm}.$$

We now show that the data of Lemma 8.13 satisfy the properties a Lorentzian \mathbb{K} -spectral triple.

Proposition 8.14. *The data $(\mathbb{K}(\Lambda, \varpi), \mathcal{V}_\Lambda \oplus \mathcal{V}_\Lambda, \mathcal{D}_\mathbb{K})$ define a Lorentzian \mathbb{K} -spectral triple, as in Definition 8.6.*

Proof. The T_ϵ are Krein isometries, since

$$(T_\epsilon e_{\lambda, \pm}, T_\epsilon e_{\lambda, \pm}) = N(\epsilon^{\rho(\lambda)})(e_{\lambda, \pm}, e_{\lambda, \pm}) = (e_{\lambda, \pm}, e_{\lambda, \pm}).$$

They satisfy $T_\epsilon^\dagger = T_{\epsilon^{-1}} = T_\epsilon^{-1}$. Thus we have $U_\epsilon^\dagger = J^\dagger T_\epsilon^\dagger = J T_\epsilon^{-1} = U_\epsilon^{-1}$. This is also a Krein isometry since both T_ϵ and J are, with $\text{Dom}(U_\epsilon) = \text{Dom}(T_\epsilon)$, since J is bounded.

The operator U_ϵ is a symmetry of the Dirac operator, namely we have

$$(8.25) \quad \mathcal{D}_{\mathbb{K}, \epsilon} := U_\epsilon^\dagger \mathcal{D}_\mathbb{K} U_\epsilon = \mathcal{D}_\mathbb{K}.$$

In fact, we have

$$(8.26) \quad T_\epsilon^\dagger \mathcal{D}_\mathbb{K} T_\epsilon e_{\lambda, \pm} = \begin{pmatrix} 0 & \epsilon^{-2\rho(\lambda)} \lambda_1 \\ \epsilon^{2\rho(\lambda)} \lambda_2 & 0 \end{pmatrix} e_{\lambda, \pm}.$$

Since we have $\lambda = A_\epsilon^{\rho(\lambda)}(\mu)$ with $\mu \in \mathcal{F}_V$, we can write the above equivalently as

$$T_\epsilon^\dagger \mathcal{D}_\mathbb{K} T_\epsilon e_{\lambda, \pm} = \begin{pmatrix} 0 & \epsilon^{-\rho(\lambda)} \mu_1 \\ \epsilon^{\rho(\lambda)} \mu_2 & 0 \end{pmatrix} e_{\lambda, \pm}.$$

Thus, we have

$$\begin{aligned} J^\dagger T_\epsilon^\dagger \mathcal{D}_\mathbb{K} T_\epsilon J e_{\lambda, \pm} &= \begin{pmatrix} 0 & \epsilon^{\rho(\lambda)} \mu_1 \\ \epsilon^{-\rho(\lambda)} \mu_2 & 0 \end{pmatrix} e_{\lambda, \pm} \\ &= \begin{pmatrix} 0 & \lambda_1 \\ \lambda_2 & 0 \end{pmatrix} e_{\lambda, \pm} = \mathcal{D}_\mathbb{K} e_{\lambda, \pm}. \end{aligned}$$

This proves property (1) of Definition 8.6.

(2) follows from Lemma 8.13 and the fact that U_ϵ is a Krein isometry, since

$$\begin{aligned} (U_\epsilon^\dagger [\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] U_\epsilon v, U_\epsilon^\dagger [\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] U_\epsilon v) &= ([\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] U_\epsilon v, [\mathcal{D}_\mathbb{K}, R_\lambda^\varpi] U_\epsilon v) \\ &= N(\lambda)(U_\epsilon v, U_\epsilon v) = N(\lambda)(v, v). \end{aligned}$$

(3) The adjoint U_ϵ^* in the associated Hilbert space inner product $\langle \cdot, \cdot \rangle$ of (8.7) satisfies $U_\epsilon^* = U_\epsilon$ since

$$\begin{aligned} U_\epsilon^* e_{\lambda, \pm} &= (\kappa U_\epsilon^\dagger \kappa) e_{\lambda, \pm} = c(U_\epsilon^\dagger) e_{\lambda, \pm} \\ &= \begin{pmatrix} c(\epsilon^{\rho(\lambda)}) & 0 \\ 0 & c(\epsilon^{-\rho(\lambda)}) \end{pmatrix} e_{J(\lambda), \pm} = \begin{pmatrix} \epsilon^{-\rho(\lambda)} & 0 \\ 0 & \epsilon^{\rho(\lambda)} \end{pmatrix} e_{J(\lambda), \pm} = U_\epsilon e_{\lambda, \pm}. \end{aligned}$$

Consider then the operator $|\mathcal{D}_\mathbb{K}^2|$ acting on the associated real Hilbert space by

$$|\mathcal{D}_\mathbb{K}^2| e_{\lambda, \pm} = \begin{pmatrix} |N(\lambda)| & 0 \\ 0 & |N(\lambda)| \end{pmatrix} e_{\lambda, \pm}.$$

We restrict $|\mathcal{D}_{\mathbb{K}}^2|$ to the orthogonal complement of the zero modes, *i.e.* on the span of the $e_{\lambda,\pm}$ with $\lambda \neq 0$. We then obtain

$$(8.27) \quad \sum_{\lambda \neq 0} |\langle U_{\epsilon} e_{\lambda,\pm}, |\mathcal{D}_{\mathbb{K}}^2 U_{\epsilon} e_{\lambda,\pm} \rangle|^{-s/2} = \sum_{\lambda \neq 0} (\epsilon^{2\rho(\lambda)} + \epsilon^{-2\rho(\lambda)})^{-s/2} |N(\lambda)|^{-s/2}.$$

This can be written equivalently as

$$(8.28) \quad \sum_{k \in \mathbb{Z}} (\epsilon^{2k} + \epsilon^{-2k})^{-s/2} \sum_{\mu \in (\Lambda \setminus \{0\})/V} |N(\mu)|^{-s/2},$$

using the unique decomposition $\lambda = A_{\epsilon}^k(\mu)$, for $k \in \mathbb{Z}$ and $\mu \in \mathcal{F}_V$, associated to the choice of the fundamental domain. Thus, we see that the finite summability condition holds. \square

Definition 8.15. *The eta function of a Lorentzian \mathbb{K} -spectral triple is the function*

$$(8.29) \quad \eta_{\mathcal{D}}(s) := \sum_n \text{sign}(\langle U e_n, \mathcal{D}^2 U e_n \rangle) |\langle U e_n, |\mathcal{D}^2 U e_n \rangle|^{-s/2},$$

where the sum is over an orthonormal basis for the complement of the zero modes of $|\mathcal{D}^2|$ in the Hilbert space $\mathcal{V}_{\mathbb{R},i}$.

The following result relates the Shimizu L -function to the Lorentz \mathbb{K} -spectral triple.

Corollary 8.16. *The eta function for the Lorentz \mathbb{K} -spectral triple of Proposition 8.14 is of the form*

$$(8.30) \quad \eta_{\mathcal{D}}(s) = L(\Lambda, V, s/2) Z_{\epsilon}(s/2),$$

where $L(\Lambda, V, s)$ is the Shimizu L -function and $Z_{\epsilon}(s/2) = \sum_{k \in \mathbb{Z}} (\epsilon^{2k} + \epsilon^{-2k})^{-s/2}$.

Proof. The argument is the same as in Proposition 8.14. We have

$$\begin{aligned} & \sum_{\lambda \neq 0} \text{sign}(\langle U_{\epsilon} e_{\lambda,\pm}, \mathcal{D}_{\mathbb{K}}^2 U_{\epsilon} e_{\lambda,\pm} \rangle) |\langle U_{\epsilon} e_{\lambda,\pm}, |\mathcal{D}_{\mathbb{K}}^2 U_{\epsilon} e_{\lambda,\pm} \rangle|^{-s/2} \\ &= \sum_{\lambda \neq 0} \text{sign}(N(\lambda)) (\epsilon^{2\rho(\lambda)} + \epsilon^{-2\rho(\lambda)})^{-s/2} |N(\lambda)|^{-s/2} \\ &= \sum_{k \in \mathbb{Z}} (\epsilon^{2k} + \epsilon^{-2k})^{-s/2} \sum_{\mu \in (\Lambda \setminus \{0\})/V} \text{sign}(N(\mu)) |N(\mu)|^{-s/2}. \end{aligned}$$

The result then follows since $L(\Lambda, V, s) = \sum_{\mu \in (\Lambda \setminus \{0\})/V} \text{sign}(N(\mu)) |N(\mu)|^{-s}$. \square

8.5. Eta function and 3-dimensional geometry. The zeta and eta functions we obtained in Proposition 8.14 and Corollary 8.16 for the Lorentzian spectral geometry are closely related to those one can obtain from the spectral geometry of the 3-dimensional solvmanifold X_{ϵ} and the signature operator on the noncommutative torus.

We have seen in §7.4 above that the Dirac operator $\not{D}_{X_{\epsilon}}$ on the 3-dimensional solvmanifold X_{ϵ} can be related to the signature operator $\not{D}_{\theta,\theta'}$ of (7.33) on the noncommutative torus $\mathbb{T}_{\Lambda,i}$. Up to a unitary equivalence, we have written in (7.36)

the operator $\tilde{D}_{\theta,\theta'}$ in terms of the operators $\tilde{D}_{\theta,\theta'}^\mu = D_\theta^\mu B_\theta$, with $\mu \in (\Lambda \setminus \{0\})/V$, defined as in (7.37).

The zeta and eta functions for the operator $\tilde{D}_{\theta,\theta'}^\mu$ have the following form.

Lemma 8.17. *The operator $\tilde{D}_{\theta,\theta'}$ has an associated zeta function of the form*

$$(8.31) \quad \zeta_{\tilde{D}_{\theta,\theta'}}(s) = 2Z_\epsilon(s/2) \sum_{\mu \in (\Lambda \setminus \{0\})/V} |N(\mu)|^{-s/2}.$$

The eta function of $\tilde{D}_{\theta,\theta'}$ vanishes due to the symmetry in the spectrum. However, the restriction $\tilde{D}_{\theta,\theta'}^+$ of $\tilde{D}_{\theta,\theta'}$ to the subspace \mathcal{H}^+ of the positive modes of the operator B_θ has a nonvanishing eta function of the form

$$(8.32) \quad \eta_{\tilde{D}_{\theta,\theta'}^+}(s) = L(\Lambda, V, s/2) Z_\epsilon(s/2) = \eta_{\mathcal{D}_\mathbb{K}}(s).$$

Proof. The operator

$$B_\theta = \begin{pmatrix} 0 & \epsilon^{-k} - i\epsilon^k \\ \epsilon^{-k} + i\epsilon^k & 0 \end{pmatrix}$$

has spectrum

$$\text{Spec}(B_\theta) = \{\pm(\epsilon^{2k} + \epsilon^{-2k})^{1/2}\}$$

which is symmetric around zero. Thus, for the zeta function we have

$$\zeta_{\tilde{D}_{\theta,\theta'}}(s) = \sum_{\mu \in (\Lambda \setminus \{0\})/V} |N(\mu)|^{-s/2} 2 \sum_{k \in \mathbb{Z}} (\epsilon^{2k} + \epsilon^{-2k})^{-s/2},$$

while the eta function vanishes.

One can restrict the spectral triple for the noncommutative torus $\mathbb{T}_{\Lambda,i}$ to the subspace \mathcal{H}^+ of the positive modes of the operator B_θ , since the action of the R_η^σ preserve this decomposition. The new Dirac operator $\tilde{D}_{\theta,\theta'}^+$ is then given by the restriction of $\tilde{D}_{\theta,\theta'}$ to \mathcal{H}^+ . It has a corresponding decomposition

$$\tilde{D}_{\theta,\theta',0}^+ = \sum_{\mu \in (\Lambda \setminus \{0\})/V} D_\theta^\mu B_\theta^+,$$

where $\text{Spec}(B_\theta^+) = \{(\epsilon^{2k} + \epsilon^{-2k})^{1/2}\}$. Thus, in this case one obtains

$$\eta_{\tilde{D}_{\theta,\theta'}^+}(s) = L(\Lambda, V, s/2) \zeta_{B_\theta^+}(s),$$

where

$$\zeta_{B_\theta^+}(s) = Z_\epsilon(s/2) = \sum_k (\epsilon^{2k} + \epsilon^{-2k})^{-s/2}.$$

□

8.6. The residue. The special value $L(\Lambda, V, 0)$ of the Shimizu L -function can be extracted from the eta function $\eta_{\tilde{p}_{\theta, \theta'}^+}(s)$ in the following way.

Corollary 8.18. *The function $\eta_{\tilde{p}_{\theta, \theta'}^+}(s)$ has a pole of order one at $s = 0$ with*

$$(8.33) \quad \text{Res}_{s=0} \eta_{\tilde{p}_{\theta, \theta'}^+}(s) = \frac{L(\Lambda, V, 0)}{\log \epsilon}.$$

Proof. Consider the function

$$Z_\epsilon(s) := \sum_{k \in \mathbb{Z}} (\epsilon^{2k} + \epsilon^{-2k})^{-s}.$$

It suffices to show that it has a simple pole at $s = 0$ with residue

$$(8.34) \quad \text{Res}_{s=0} Z_\epsilon(s) = \frac{1}{\log \epsilon}.$$

One writes

$$\Gamma(s) Z_\epsilon(s) = \int_0^\infty g_\epsilon(t) t^{s-1} dt,$$

where

$$(8.35) \quad g_\epsilon(t) = \left(\sum_{k \in \mathbb{Z}} e^{-(\epsilon^{2k} + \epsilon^{-2k})t} \right)$$

for $t > 0$. The function $g_\epsilon(t)$ satisfies

$$g_\epsilon(t) = -e^{-2t} + 2h_\epsilon(t) - 2 \sum_{k=0}^\infty e^{-\epsilon^{2k}t} (1 - e^{-\epsilon^{-2k}t}),$$

where

$$h_\epsilon(t) = \sum_{k=0}^\infty e^{-\epsilon^{2k}t}.$$

We can estimate $-e^{-2t} = -1 + O(t)$ when $t \rightarrow 0$ and $(1 - e^{-\epsilon^{-2k}t}) = O(\epsilon^{-2k}t)$, uniformly. Notice that

$$h_\epsilon(t) - h_\epsilon(\epsilon^2 t) = e^{-t} = \sum_{r=0}^\infty \frac{(-1)^r}{r!} t^r$$

hence

$$(8.36) \quad h_\epsilon(t) = \frac{1}{2 \log \epsilon} \log(1/t) + C - \sum_{r=0}^\infty \frac{(-1)^r}{r! (\epsilon^{2r} - 1)} t^r.$$

Thus, the function $\Gamma(s) Z_\epsilon(s)$ has a double pole at $s = 0$ and simple poles at $s \in \mathbb{Z}_{<0}$. Thus, the function $Z_\epsilon(s)$ has a simple pole at zero with residue $1/\log \epsilon$. \square

REFERENCES

- [1] M.F. Atiyah, *Elliptic operators, discrete groups and von Neumann algebras*. Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974), pp. 43–72. Asterisque, No. 32-33, Soc. Math. France, Paris, 1976.
- [2] M.F. Atiyah, H. Donnelly, I.M. Singer, *Eta invariants, signature defects of cusps, and values of L-functions*, Annals of Mathematics 118 (1983) 131–177.
- [3] M.F. Atiyah, V.K. Patodi, I.M. Singer, *Spectral asymmetry and Riemannian geometry. III*. Math. Proc. Cambridge Philos. Soc. 79 (1976), no. 1, 71–99.
- [4] P. Baum, A. Connes, *Chern character for discrete groups*. A fete of topology, 163–232, Academic Press, Boston, MA, 1988.
- [5] J. Bellissard, *Noncommutative geometry and quantum Hall effect*. Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994), 1238–1246, Birkhäuser, 1995.
- [6] J. Bellissard, *Gap labelling theorems for Schrödinger operators*, From number theory to physics (Les Houches, 1989), 538–630, Springer, Berlin, 1992.
- [7] J. Bognár, *Indefinite inner product spaces*, Springer, 1974.
- [8] J. Chabert, S. Echterhoff, *Permanence properties of the Baum–Connes conjecture*, Doc. Math. 6 (2001) 127–183.
- [9] A. Connes, *C^* -algèbres et géométrie différentielle*. C. R. Acad. Sci. Paris Sér. A-B 290 (1980), no. 13, A599–A604.
- [10] A. Connes, *An analogue of the Thom isomorphism for crossed products of a C^* -algebra by an action of \mathbb{R}* . Adv. in Math. 39 (1981), no. 1, 31–55.
- [11] A. Connes, *Geometry from the spectral point of view*. Lett. Math. Phys. 34 (1995), no. 3, 203–238.
- [12] A. Connes, G. Landi, *Noncommutative manifolds, the instanton algebra and isospectral deformations*. Comm. Math. Phys. 221 (2001), no. 1, 141–159.
- [13] S. Echterhoff, W. Lück, N.C. Phillips, S. Walters, *The structure of crossed products of irrational rotation algebras by finite subgroups of $SL_2(\mathbb{Z})$* , preprint SFB 478, Heft 435, 2006.
- [14] E. Getzler, *The odd Chern character in cyclic homology and spectral flow*. Topology 32 (1993), no. 3, 489–507.
- [15] F. Hirzebruch, *Hilbert modular surfaces*. Enseignement Math. (2) 19 (1973), 183–281.
- [16] D.G. Hofstadter, *Energy levels and wave functions of Bloch electrons in rational or irrational magnetic field*. Phys. Rev. B14 (1976) 2239–2249.
- [17] Yu.I. Manin, *Real multiplication and noncommutative geometry (ein Alterstraum)*. The legacy of Niels Henrik Abel, 685–727, Springer, Berlin, 2004.
- [18] M. Marcolli, V. Mathai, *Twisted index theory on good orbifolds, I: noncommutative Bloch theory*, Commun. Contemp. Math. Vol1 (1999) N.4, 553–587.
- [19] M. Marcolli, V. Mathai, *Twisted index theory on good orbifolds, II: fractional quantum numbers*, Commun. Math. Phys. 217 (2001) 55–87.
- [20] M. Marcolli, V. Mathai, *Towards the fractional quantum Hall effect, a noncommutative geometry perspective*, Noncommutative Geometry and Number Theory, pp.235–262. Vieweg Verlag, 2006.
- [21] R. Matthes, O. Richter, G. Rudolph, *Spectral triples and differential calculi related to the Kronecker foliation*. J. Geom. Phys. 46 (2003), no. 1, 48–73.
- [22] A. Mishchenko, A. Fomenko, *The index of elliptic operators over C^* -algebras*. Math. USSR Izv. 15 (1980) 87–112.
- [23] W. Müller, *Signature defects of cusps of Hilbert modular varieties and values of L-series at $s = 1$* . J. Differential Geom. 20 (1984), no. 1, 55–119.
- [24] J.A. Packer, I. Raeburn, *On the structure of twisted group C^* -algebras*, Trans. AMS 334 (1992) 685–718.
- [25] A. Rennie, J. Varilly, *Reconstruction of manifolds in noncommutative geometry*, arXiv:math/0610418.

- [26] M.A. Rieffel, *C^* -algebras associated with irrational rotations*. Pacific J. Math. 93 (1981), no. 2, 415–429.
- [27] J. Rosenberg, *C^* -algebras, positive scalar curvature, and the Novikov conjecture III*, Topology 25 (1986) 319–336.
- [28] A. Strohmaier, *On noncommutative and pseudo-Riemannian geometry*. J. Geom. Phys. 56 (2006), no. 2, 175–195.
- [29] T. Sunada, *A discrete analogue of periodic magnetic Schrödinger operators*. Geometry of the spectrum (Seattle, WA, 1993), 283–299, Contemp. Math., 173, Amer. Math. Soc., Providence, RI, 1994.

MAX-PLANCK INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, BONN, D-53111 GERMANY
E-mail address: marcolli@mpim-bonn.mpg.de